arXiv:1409.3729v2 [math.AG] 14 Jul 2015

LAURENT PHENOMENON FOR LANDAU–GINZBURG MODELS OF COMPLETE INTERSECTIONS IN GRASSMANNIANS OF PLANES

VICTOR PRZYJALKOWSKI, CONSTANTIN SHRAMOV

Abstract. In a spirit of Givental's constructions Batyrev, Ciocan-Fontanine, Kim, and van Straten suggested Landau–Ginzburg models for smooth Fano complete intersections in Grassmannians and partial flag varieties as certain complete intersections in complex tori equipped with special functions called superpotentials. We provide a particular algorithm for constructing birational isomorphisms of these models for complete intersections in Grassmannians of planes with complex tori. In this case the superpotentials are given by Laurent polynomials. We study Givental's integrals for Landau–Ginzburg models suggested by Batyrev, Ciocan-Fontanine, Kim, and van Straten and show that they are periods for pencils of fibers of maps provided by Laurent polynomials we obtain. The algorithm we provide after minor modifications can be applied in a more general context. In the appendix written by Thomas Prince one more algorithm is presented.

Contents

1. Introduction 1
2. Toric Landau–Ginzburg models 4
3. Complete intersections in smooth toric varieties 6
4. Complete intersections in Grassmannians 10
5. Main theorem 13
6. Horizontal blocks 16
7. Mixed blocks 31
8. Vertical blocks 37
9. Proof of the main theorem 41
10. Periods 43
11. Hyperplane sections 47
12. Examples 50
13. Discussion 60
Appendix A. Laurent polynomial mirrors for Grassmannians of planes, by Thomas Prince 62
References 66

1. Introduction

Mirror Symmetry declares duality between algebraic and symplectic geometries of different varieties. Starting from duality between Calabi–Yau varieties it was extended to Fano varieties (see e. g. [Ko94]). In this case the dual object to a Fano variety is called a Landau–Ginzburg model. Its definition varies depending on a version of Mirror Symmetry

This work is supported by the Russian Science Foundation under grant 14-50-00005.

conjectures. In what follows mirror partners for us are smooth Fano varieties and their toric Landau–Ginzburg models.

The challenge for Mirror Symmetry is to find mirror partners for given varieties and, via studying them, explore (or check or at least guess) geometry of the initial varieties. In the paper we mostly focus on the problem of finding Landau–Ginzburg models. Historically constructions of Landau–Ginzburg models, initiated by Givental ([Gi97b]), Eguchi, Hori and Xiong ([EHX97]), Batyrev ([Ba97]), Batyrev, Ciocan-Fontanine, Kim and van Straten ([BCFKS98]), Hori and Vafa ([HV00]), and other people, were based on a toric approach. The idea is, given a smooth toric Fano variety or a variety having a (nice) toric degeneration, to construct a Laurent polynomial whose support is a fan polytope of a toric variety or a toric degeneration. This idea is initiated by Batyrev–Borisov approach to treat mirror duality for a toric variety as a classical duality of toric varieties corresponding to dual polytopes. The constructed Laurent polynomial gives a map from a complex torus to an affine line and, thus, defines a Landau–Ginzburg model. We do not discuss methods of finding appropriate Laurent polynomial in a general case; see [Ba97], [Prz13], [CCGGK12] for details.

In [Gi97b] (see discussion after Corollary 0.4 therein) Givental suggested an approach to writing down Landau–Ginzburg models for complete intersections in toric varieties or varieties having nice (say, terminal Gorenstein) toric degenerations (see also [HV00, §7.2]). This approach assumes an existence of a *nef-partition* — a subdivision of a set of rays of a fan of a toric variety into disjoint subsets such that the sum of invariant divisors corresponding to each subset is linearly equivalent to a hypersurface defining the complete intersection.

In some cases this construction is described in details (see e. g. [BCFKS98], [HV00, §7.2], [DHKLP]). A priori the output of the construction is a quasiaffine variety with a complex-valued function (called *superpotential*) on it. However in many cases such quasiaffine varieties are birational to tori and thus the Landau–Ginzburg models are toric (which is very useful for studying them), see, for instance, [Prz10]. We can't prove this phenomenon in the general case: there is no approach to construct (or even prove an existence of) a nef-partition. In some cases, however, the nef-partition is described. Landau–Ginzburg models for Grassmannians themselves were suggested in [EHX97, B25] as Laurent polynomials, i. e. functions on tori already. In [BCFKS97] and [BCFKS98] the description for Landau–Ginzburg models for complete intersections in Grassmannians and partial flag varieties as complete intersections in tori are given. However it is not clear if they themselves are birational to tori (cf. [Prz13, Problem 16]). We prove the phenomenon in the case of complete intersections of Grassmannians of planes.

A basic and the most important property of Landau–Ginzburg models from the point of view discussed above is a *period condition*. It relates (some of) the periods of a Landau–Ginzburg model for a Fano variety with its *I-series*, a generating series for one-pointed Gromov–Witten invariants. These series for Grassmannians are found in [BCFK03]. Moreover, Quantum Lefschetz Theorem (see for instance [Gi97b, Theorem 0.1], [Ki00], [Lee01]) relates I-series of a Fano variety and a complete intersection therein with nef anticanonical class. Thus this series is known for complete intersections in Grassmannians as well. Periods for these complete intersections are proven to be related with their I-series in [BCFKS98] modulo the assumption that this holds for Grassmannians themselves. This assumption is proven for Grassmannians of planes

in [BCFK03, Proposition 3.5] and for all Grassmannians in [MR13]. Thus the period condition holds for Landau–Ginzburg models for smooth Fano complete intersections in Grassmannians suggested in [BCFKS97].

A very weak Landau–Ginzburg model is a Laurent polynomial for which the period condition holds. The stronger notion, a notion of *toric Landau–Ginzburg model*, requires two more conditions. *A Calabi–Yau condition* states an existence of a relative compactification of the family that is a (non-compact) Calabi–Yau variety. *A toric condition* states that a Newton polytope of the Laurent polynomial is a fan polytope of a toric degeneration of the Fano variety.

Theorem 1.1 (Corollary 10.6)**.** *Any smooth Fano complete intersection in a Grassmannian of planes has a very weak Landau–Ginzburg model.*

The particular form of this very weak Landau–Ginzburg model can be derived from Theorem 5.3.

Conjecture 1.2. *The assertion of Theorem 1.1 holds for complete intersections in any Grassmannian or, more generally, partial flag variety.*

This conjecture is proved by methods different from ones used in this paper for complete intersections in Grassmannians and for a big class of complete intersections in partial flag varieties in [DH15] and for complete intersections in Grassmannians in [PSh15b].

Theorem 1.1 is one more evidence for the following conjecture, that may be regarded as a strong version of Mirror Symmetry of variations of Hodge structures conjecture, see [Prz13, Conjecture 38].

Conjecture 1.3. *Any smooth Fano variety has a toric Landau–Ginzburg model.*

In particular, we have the following weaker conjectures.

Conjecture 1.4. *Any smooth Fano variety has a toric degeneration.*

Conjecture 1.5. *Any smooth Fano variety has a rational Landau–Ginzburg model of the same dimension.*

The paper is organized as follows. In Section 2 we give definitions of toric Landau–Ginzburg models and I-series for Fano varieties. In Section 3 we give definitions of Givental's period integrals and Givental's Landau–Ginzburg models for complete intersections in smooth toric varieties. They are defined via nef-partitions and relations between rays of a fan defining the toric variety. Also, in Section 3 we show how changing variables one can simplify Givental's integrals and Landau–Ginzburg models and get rid of the part depending on the relations.

Givental's approach can be applied in a more general case. That is one can define Landau–Ginzburg models and period integrals of smooth Fano complete intersections in Fano varieties having "good", say terminal Gorenstein, toric degenerations. To do this one applies the Givental's method to a certain complete intersection in a crepant resolution of the toric degeneration of the latter Fano variety and then taking a specializations of some parameters. Following [BCFKS98] we show how it works for Grassmannians of planes in Section 4. In Section 5 we reformulate notions given in Section 4 in a more abstract way suitable for the proof of our main assertions, and formulate Theorem 5.3 that is a more technical counterpart of Theorem 1.1. Sections 6–8 are pretty big and contain our

main technical lemmas needed for the proof of Theorem 5.3. The proof itself is given in Section 9.

Section 10 is devoted to further simplifications of Givental's integrals. Theorem 5.3 states that there is a series of changes of variables allowing one to birationally present Landau–Ginzburg models for complete intersections in Grassmannians of planes as Laurent polynomials. In Section 10 we check that these changes of variables (and also ones for complete intersections in projective spaces) agree with changing Givental's integrals. This shows that Givental's integrals are indeed periods; they can be easily computed as constant term series. As a corollary one gets the fact that complete intersections in Grassmannians of planes have very weak Landau–Ginzburg models.

In Section 11 we write down explicit formulas that are obtained in the proof of Theorem 5.3 for Fano intersections of Grassmannians of planes with several hyperplanes. Section 12 provides a series of examples worked out as an application of Theorem 5.3 in some other interesting cases.

Those who have a certain amount of combinatorial courage can apply the approaches described in our paper in more general cases, say, for complete intersections in arbitrary Grassmannians or even in partial flag varieties. We discuss this in Section 13. We also discuss how one can study very weak toric Landau–Ginzburg models given in the proof of Theorem 5.3 in a deeper way.

Finally in the Appendix written by Thomas Prince another algorithm to transform Landau–Ginzburg models suggested in [BCFKS97] to very weak Landau–Ginzburg models is presented. It uses a completely different approach inspired by [CKP14] and gives a shorter way to obtain Theorem 5.3.

Notation and conventions. Everything is over $\mathbb{C}$. We use (co-)homology groups with integral coefficients and denote $H^*(X,\mathbb{Z})$ by $H^*(X)$ and $H_*(X,\mathbb{Z})$ by $H_*(X)$. Given two integers n_1 and n_2, we denote the set $\{i \mid n_1 \leqslant i \leqslant n_2\}$ by $[n_1,n_2]$. Calabi–Yau varieties in this paper are projective varieties with trivial canonical class. We often use the same notation for a (Cartier) divisor on a variety X and its class in $\mathrm{Pic}\,(X)$. When we speak about hyperplane or hypersurface sections of a Grassmannian we mean hyperplane or hypersurface sections in its Plücker embedding.

Acknowledgments. The authors are very grateful to B. Kim and I. Ciocan-Fontanine for explanations of Givental's integrals and for many useful remarks, and also to A. Harder, A. Fonarev, S. Gorchinskiy, and A. Kuznetsov for inspiring discussions.

2. Toric Landau–Ginzburg models

In this section we define the main objects of our considerations — toric Landau–Ginzburg models. For more details and examples see [Prz13] and references therein.

Let X be a smooth Fano variety of dimension N and Picard number ρ. Choose a basis

$$\{H_1,\dots,H_\rho\}$$

in $H^2(X)$ so that for any $i \in [1,\rho]$ and any curve β in the Kähler cone K of X one has $H_i \cdot \beta \geqslant 0$. Introduce formal variables q^{τ_i}, $i \in [1,\rho]$ and denote $q_i = q^{\tau_i}$. For any $\beta \in H_2(X)$ denote

$$q^\beta = q^{\sum \tau_i (H_i \cdot \beta)}.$$

Consider the Novikov ring $\mathbb{C}_q$, i. e. a group ring for $H_2(X)$. We treat it as a ring of polynomials over $\mathbb{C}$ in formal variables q^β, with relations

$$q^{\beta_1} q^{\beta_2} = q^{\beta_1+\beta_2}.$$

Note that for any $\beta \in K$ the monomial q^β has non-negative degrees in q_i.

Let the number

$$\langle \tau_a \gamma \rangle_\beta, \quad a \in \mathbb{Z}_{\geq 0},\ \gamma \in H^*(X),\ \beta \in K,$$

be a one-pointed Gromov–Witten invariant with descendants for X, see [Ma99, VI-2.1]. Let $\mathbf{1}$ be the fundamental class of X. The series

$$I_0^X(q_1, \dots, q_\rho) = 1 + \sum_{\beta \in K} \langle \tau_{-K_X \cdot \beta - 2} \mathbf{1} \rangle_{-K_X \cdot \beta} \cdot q^\beta$$

is called *a constant term of I-series* (or *a constant term of Givental's J-series*) for X and the series

$$\widetilde{I}_0^X(q_1, \dots, q_\rho) = 1 + \sum_{\beta \in K} (-K_X \cdot \beta)! \langle \tau_{-K_X \cdot \beta - 2} \mathbf{1} \rangle_{-K_X \cdot \beta} \cdot q^\beta$$

is called *a constant term of regularized I-series* for X. Given a divisor $H = \sum \alpha_i H_i$ one can restrict these series to a direction corresponding to this divisor setting $\tau_i = \alpha_i \tau$ and $t = q^\tau$. Thus one can define a *restriction of a constant term of regularized I-series to anticanonical direction*; it has the form

$$\widetilde{I}^X(t) = 1 + a_1 t + a_2 t^2 + \dots.$$

Definition 2.1 (see [Prz13, §6])**.** *A toric Landau–Ginzburg model* of X is a Laurent polynomial $f \in \mathbb{C}[x_1^{\pm 1}, \dots, x_N^{\pm 1}]$ which satisfies:

Period condition: The constant term of f^i equals a_i for any i.

Calabi–Yau condition: There exists a relative compactification of a family

$$f \colon (\mathbb{C}^*)^N \to \mathbb{C}$$

whose total space is a (non-compact) smooth Calabi–Yau variety $LG(X)$. Such compactification is called *a Calabi–Yau compactification*.

Toric condition: There is a degeneration $X \rightsquigarrow T$ to a toric variety T whose fan polytope (i. e. the convex hull of generators of its rays) coincides with Newton polytope (i. e. the convex hull of the support) of f.

Remark 2.2. The period condition is a numerical expression of coincidence of constant term of regularized I-series and a period of the family provided by f, see Remark 10.2 and Theorem 10.3.

Let us remind that the Laurent polynomials for which the period condition is satisfied are called very weak Landau–Ginzburg models; ones for which in addition a Calabi–Yau condition holds are called *weak Landau–Ginzburg models*.

Toric Landau–Ginzburg models are known for Fano threefolds (see [Prz13], [ILP13], and [DHKLP]) and complete intersections in projective spaces ([ILP13]); some other partial results are also known.

There are two usual ways to find toric Landau–Ginzburg models. The first way is to find birational transformations of known suggestions for Landau–Ginzburg models to make their total spaces tori. In this case their superpotentials in toric coordinates are Laurent polynomials. After this one can try to prove the three conditions for being a toric

Landau–Ginzburg model. An important case of this approach is the following. Given a Fano variety one can sometimes describe it (or its "good" degeneration) as a complete intersection in a toric variety. Then one can try to find a relative birational isomorphism of Givental's type Landau–Ginzburg models (see Definition 3.3) with a torus. An example of this approach can be found in [Prz10] and [CCGGK12].

The second way is to find a toric degenerations of X. Given this degeneration one has a Newton polytope of its possible toric Landau–Ginzburg model and so can try to find its particular coefficients. One can look at [DHKLP] for an example of this approach.

In this paper we apply the first approach to find candidates for toric Landau–Ginzburg models for complete intersections in Grassmannians using Givental suggestions of Landau–Ginzburg models. We conjecture that Laurent polynomials we get are toric Landau–Ginzburg models.

3. Complete intersections in smooth toric varieties

In this section we describe Givental's construction of Landau–Ginzburg models and period integrals for complete intersections in toric varieties given in [Gi97b] (see discussion after Corollary 0.4 therein). That is, we describe their very weak Landau–Ginzburg models and discuss their periods. Further in Section 4 we literally repeat these considerations for complete intersections in singular toric varieties (which are terminal Gorenstein degenerations of Grassmannians).

Let X be a factorial N-dimensional toric Fano variety of Picard rank ρ corresponding to a fan Σ_X in a lattice $\mathcal{N} \cong \mathbb{Z}^N$. Let $D_1, \dots, D_{N+\rho}$ be its prime invariant divisors. Let $Y_1, \dots, Y_l$ be ample divisors in X cutting out a smooth Fano complete intersection

$$Y = Y_1 \cap \ldots \cap Y_l.$$

Put $Y_0 = -K_X - Y_1 - \ldots - Y_l$. Choose a basis

$$\{H_1, \dots, H_\rho\} \subset H^2(X)$$

so that for any $i \in [1, \rho]$ and any curve β in the Kähler cone K of X one has $H_i \cdot \beta \geqslant 0$. Introduce variables $q_1, \dots, q_\rho$ as in Section 2.

The following theorem is a particular case of Quantum Lefschetz hyperplane theorem, see [Gi97b, Theorem 0.1].

Theorem 3.1. *Suppose that* $\dim(Y) \geqslant 3$. *Then the constant term of regularized I-series for Y is given by*

$$\widetilde{I}_0^Y(q_1, \dots, q_\rho) = \exp\big(\mu(q)\big) \cdot \sum_{\beta \in K \cap H_2(X)} q^\beta \frac{\prod_{i=0}^{l} |\beta \cdot Y_i|!}{\prod_{j=1}^{N+\rho} |\beta \cdot D_j|!^{\frac{\beta \cdot D_j}{|\beta \cdot D_j|}}} \tag{3.1}$$

where $\mu(q)$ is a correction term linear in q_i (in particular it is trivial in the higher index case). For $\dim(Y) = 2$ *the same formula holds after replacing $H^2(Y)$ in the definition of $\widetilde{I}_0^Y$ given in Section 2 by the restriction of $H^2(X)$ to Y.*

Remark 3.2. Note that the summands of the series (3.1) have non-negative degrees in q_i.

Now we describe Givental's construction of a dual Landau–Ginzburg model of Y and compute its periods. Introduce N formal variables $u_1, \dots, u_{N+\rho}$ corresponding to divisors $D_1, \dots, D_{N+\rho}$.

Let $\mathcal{M} = \mathcal{N}^\vee$, and let $\mathcal{D} \cong \mathbb{Z}^{N+\rho}$ be a lattice with a basis $\{D_1, \dots, D_{N+\rho}\}$ (so that one has a natural identification $\mathcal{D} \cong \mathcal{D}^\vee$). By [CLS11, Theorem 4.2.1] one has an exact sequence

$$0 \to \mathcal{M} \to \mathcal{D} \to A_{N-1}(X) = \operatorname{Pic}(X) \cong \mathbb{Z}^\rho \to 0.$$

We use factoriality of X here to identify the class group $A_{N-1}(X)$ and the Picard group $\operatorname{Pic}(X)$. Dualizing this exact sequence, we obtain an exact sequence

$$0 \to \operatorname{Pic}(X)^\vee \to \mathcal{D} \to \mathcal{N} \to 0. \tag{3.2}$$

Thus $\operatorname{Pic}(X)^\vee$ can be identified with the lattice of relations on primitive vectors on the rays of Σ_X considered as Laurent monomials in variables u_i. On the other hand, as the basis in $\operatorname{Pic}(X)$ is chosen we can identify $\operatorname{Pic}(X)^\vee$ and $\operatorname{Pic}(X) = H^2(X)$. Hence we can choose a basis in the lattice of relations on primitive vectors on the rays of Σ_X corresponding to $\{H_i\}$ and, thus, to $\{q_i\}$. We denote these relations by R_i, and interpret them as monomials in the variables $u_1, \dots, u_{N+\rho}$. We also denote by D_i the images of $D_i \in \mathcal{D}$ in $\operatorname{Pic} X$.

Choose a nef-partition, i. e. a partition of the set $[1, N+\rho]$ into sets $E_0, \dots, E_l$ such that for any $i \in [1, l]$ the divisor $\sum_{j \in E_i} D_j$ is linearly equivalent to Y_i (which also implies that the divisor $\sum_{j \in E_0} D_j$ is linearly equivalent to Y_0).

The following definition is well-known (see discussion after Corollary 0.4 in [Gi97b], and also [HV00, §7.2]).

Definition 3.3. *Givental's Landau–Ginzburg model* for Y is a variety $LG_0(Y)$ in a torus

$$T = \operatorname{Spec} \mathbb{C}_q[u_1^{\pm 1}, \dots, u_{N+\rho}^{\pm 1}]$$

given by equations

$$R_i = q_i, \ i \in [1, \rho], \tag{3.3}$$

and

$$\left(\sum_{s \in E_j} u_s \right) = 1, \ j \in [1, l], \tag{3.4}$$

with a function $w = \sum_{s \in E_0} u_s$ called *superpotential*. Given a symplectic form ω with $[\omega] \sim \sum \omega_i H_i$, where $[\omega]$ is the class in $\operatorname{Pic}(Y)$ corresponding to ω, define *the Givental's Landau–Ginzburg model $LG(Y, \omega)$ associated to ω* specializing $q_i = \exp(\omega_i)$. If ω is an anticanonical form ω_Y, i. e. one has $[\omega] = -K_Y$, we say for simplicity that $LG(Y) = LG(Y, \omega)$ is an anticanonical Givental's Landau–Ginzburg model for Y instead of saying that $LG(Y, \omega_Y)$ is a Givental's Landau–Ginzburg model for (Y, ω_Y).

Remark 3.4. The superpotential of Givental's Landau–Ginzburg models can be defined as $w' = u_1 + \ldots + u_{N+\rho}$. However we don't make a distinction between two superpotentials w and w' as $w' = w + l$, so both these functions define the same family over $\mathbb{C}_q$.

Given variables $x_1, \dots, x_r$, define a *standard logarithmic form in these variables* as the form

$$\Omega(x_1, \dots, x_r) = \frac{1}{(2\pi i)^r} \frac{dx_1}{x_1} \wedge \ldots \wedge \frac{dx_r}{x_r}. \tag{3.5}$$

The following definition is well-known (see discussion after Corollary 0.4 in [Gi97b], and also [Gi97a]).

Definition 3.5. Fix $N+\rho$ real positive numbers $\varepsilon_1,\dots,\varepsilon_{N+\rho}$ and define an $(N+\rho)$-cycle

$$\delta=\{|u_i=\varepsilon_i|\}\subset\mathbb{C}[u_1^{\pm1},\dots,u_{N+\rho}^{\pm1}].$$

Givental's integral for Y or $LG_0(Y)$ is an integral

$$I_Y^0=\int\limits_{\delta}\frac{\Omega(u_1,\dots,u_{N+\rho})}{\prod_{i=1}^{\rho}(1-\frac{q_i}{R_i})\cdot\prod_{j=0}^{l}\left(1-\left(\sum_{s\in E_j}u_s\right)\right)}\in\mathbb{C}[[q_1,\dots,q_\rho]]. \tag{3.6}$$

Given a form $\omega=\sum\alpha_iH_i$ one can *specialize Givental's integral to the direction* ω putting $q_i=t^{\alpha_i}$ in the integral (3.6). We denote the result of specialization by $I_{(Y,\omega)}$, and we put $I_{(Y,\omega)}=I_Y$ if $[\omega]=-K_Y$.

Remark 3.6. The integral (3.6) does not depend on numbers ε_i provided they are small enough.

Remark 3.7. The integral (3.6) is defined up to a sign as we do not specify an order of variables.

The following assertion is well-known to experts (see [Gi97b, Theorem 0.1], and also discussion after Corollary 0.4 in [Gi97b]).

Theorem 3.8. *One has*

$$\widetilde{I}_0^Y=I_Y^0.$$

The recipe for Givental's Landau–Ginzburg model and integral can be written down in another, more simple, way. That is, we make suitable monomial change of variables $u_1,\dots,u_{N+\rho}$ an get rid of some of them using equations (3.3). More precisely, as $\mathcal{N}$ is a free group, using the exact sequence (3.2) one obtains an isomorphism

$$\mathcal{D}\cong\operatorname{Pic}X^\vee\oplus\mathcal{N}.$$

Thus one can find a monomial change of variables $u_1,\dots,u_{N+\rho}$ to some new variables $x_1,\dots,x_N,y_1,\dots,y_\rho$, so that

$$u_i=\widetilde{X}_i(x_1,\dots,x_N,y_1,\dots,y_\rho,q_1,\dots,q_\rho)$$

such that for any $i\in[1,\rho]$ one has

$$\frac{R_i(u_1,\dots,u_{N+\rho})}{q_i}=\frac{1}{y_i}.$$

Put

$$X_i=\widetilde{X}_i(x_1,\dots,x_N,1,\dots,1,q_1,\dots,q_\rho).$$

Then $LG(Y)$ is given in the torus $\operatorname{Spec}\mathbb{C}_q[x_1^{\pm1},\dots,x_N^{\pm1}]$ by equations

$$\left(\sum_{s\in E_j}\alpha_sX_s\right)=1,\ j\in[1,l],$$

with superpotential $w=\sum_{s\in E_0}\alpha_sX_s$, where $\alpha_i=\prod q_j^{r_{i,j}}$ for some integers $r_{i,j}$.

Let us mention that given a Laurent monomial U_i in variables u_j, $j\in[1,N+\rho]$, that does not depend on a variable u_i one has

$$\Omega(u_1,\dots,u_i^{\pm1}\cdot U_i,\dots,u_{N+\rho})=\pm\Omega(u_1,\dots,u_i,\dots,u_{N+\rho}). \tag{3.7}$$

This means that

$$I_Y^0 = \int_{\delta'} \frac{\pm\Omega(y_1,\dots,y_\rho)\wedge\Omega(x_1,\dots,x_N)}{\prod_{i=1}^{\rho}(1-y_i)\prod_{j=0}^{l}\left(1-\left(\sum_{s\in E_j}\alpha_s\widetilde{X}_s\right)\right)} \tag{3.8}$$

for some $(N+\rho)$-cycle δ'.

Consider an integral

$$\int_\sigma \frac{dU}{U}\wedge\Omega_0$$

for some form Ω_0 and a cycle $\sigma=\sigma'\cap\{|U|=\varepsilon\}$ for some cycle $\sigma'\subset\{U=0\}$. It is well known that (see, for instance, [ATY85, Theorem 1.1]) that

$$\frac{1}{2\pi i}\int_\sigma \frac{dU}{U}\wedge\Omega_0 = \int_{\sigma'}\Omega_0|_{U=0}$$

if both integrals are well defined (in particular the form Ω_0 does not have a pole along $\{U=0\}$).

We denote

$$\Omega_0|_{U=0} = \operatorname{Res}_U\left(\frac{dU}{U}\wedge\Omega_0\right).$$

Taking residues of the form on the right hand side of (3.8) with respect to y_i one gets

$$I_Y^0 = \int_{\delta''} \frac{\pm\Omega(x_1,\dots,x_N)}{\prod_{j=0}^{l}\left(1-\left(\sum_{s\in E_j}\alpha_s X_s\right)\right)}$$

for some N-cycle δ''.

Moreover, one can introduce a new parameter t and scale $u_i\to tu_i$ for $i\in E_0$. One can check that after a change of coordinates $q_i=1$ the initial integral restricts to the integral

$$\int_{\delta_1} \frac{\pm\Omega(x_1,\dots,x_N)}{\prod_{j=1}^{l}\left(1-\left(\sum_{s\in E_j}X_s\right)\right)\cdot\left(1-t\left(\sum_{i\in E_0}X_i\right)\right)} = I_Y \tag{3.9}$$

for some N-cycle δ_1 homologous to a cycle

$$\delta_1^0 = \{|x_i|=\varepsilon_i \mid i\in[1,N]\}.$$

The same specialization defines the anticanonical Givental's Landau–Ginzburg model of Y, which is given by equations

$$\left(\sum_{s\in E_j} X_s\right) = 1, \quad j\in[1,l],$$

with superpotential $w=\sum_{s\in E_0}X_s$.

Consider a non-toric variety X that has a small (that is, terminal Gorenstein) toric degeneration T. Let Y be a Fano complete intersection in X. Consider a nef-partition for the set of rays of the fan of T corresponding to (degenerations of) hypersurfaces cutting out Y. Let $LG(Y)$ be a result of applying the procedure discussed above for Givental's integral defined for T and the nef-partition in the same way as in the case of complete intersections in toric varieties. Batyrev in [Ba97] suggested $LG(Y)$ as a Landau–Ginzburg model for Y. Moreover, at least in some cases Givental's integral and Landau–Ginzburg model (associated to anticanonical class) can be simplified further by making birational changes of variables and taking residues. Thus Givental's Landau–Ginzburg models give

very weak ones after such transformations. In Section 10 we demonstrate both of these ideas for complete intersections in projective spaces or Grassmannians of planes.

4. Complete intersections in Grassmannians

The picture described in Section 3 can be generalized to complete intersections in Grassmannians. The difference is that Grassmannians are not toric. However they have *small toric degenerations*, i. e. degenerations to terminal Gorenstein toric varieties, see [St93]. The mirror construction for complete intersections in Grassmannians can be derived from crepant resolutions of these degenerations. In this section we describe some constructions from [BCFKS97] and [BCFKS98] for a Grassmannian $G = \mathrm{Gr}(n, k+n)$.

Fix two integers n and k such that $n, k \geqslant 2$. We define a quiver $\mathcal{Q}_0$ as a set of vertices

$$\mathrm{Ver}(\mathcal{Q}_0) = \{(i,j) \mid i \in [1,k], j \in [1,n]\} \cup \{(0,1), (k,n+1)\}$$

and a set of arrows $\mathrm{Ar}(\mathcal{Q}_0)$ described as follows. All arrows are either *vertical* or *horizontal*. For any $i \in [1,k-1]$ and any $j \in [1,n]$ there is one vertical arrow $\langle (i,j) \to (i+1,j) \rangle$ that goes from the vertex (i,j) down to the vertex $(i+1,j)$. For any $i \in [1,k]$ and any $j \in [1,n-1]$ there is one horizontal arrow $\langle (i,j) \to (i,j+1) \rangle$ that goes from the vertex (i,j) to the right to the vertex $(i,j+1)$. We also add an extra vertical arrow $\langle (0,1) \to (1,1) \rangle$ and an extra horizontal arrow $\langle (k,n) \to (k,n+1) \rangle$ to $\mathrm{Ar}(\mathcal{Q}_0)$, see Figure 1.



Figure 1. Quiver $\mathcal{Q}_0$ for the Grassmannian $\mathrm{Gr}(2,5)$

Now we describe a toric degeneration $P = P(n, k+n)$ of G in its Plücker embedding. The arrows of $\mathcal{Q}_0$ correspond to rays of a fan Σ_P of P, so we identify them; relations for the primitive vectors on the rays of Σ_P correspond to cycles in $\mathcal{Q}_0$ if we identify vertices $(0,1)$ and $(k,n+1)$. The cones of Σ_P of dimension at least 2 are cones over faces of a convex hull of generators of rays of Σ_P. A degeneration P is a Fano toric variety corresponding to Σ_P.

The variety P is not smooth but terminal Gorenstein. It admits (some) crepant resolution that we denote by $\widetilde{P}$. All relations on rays of P (or $\widetilde{P}$) are combinations of basic

ones described as follows. For any $i \in [1, k-1]$ and $j \in [1, n-1]$ we have a *box relation*

$$\langle (i,j) \to (i+1,j) \rangle + \langle (i+1,j) \to (i+1,j+1) \rangle = \\ = \langle (i,j) \to (i,j+1) \rangle + \langle (i,j+1) \to (i+1,j+1) \rangle;$$

besides that, we have one *roof relation*

$$0 = \langle (0,1) \to (1,1) \rangle + \langle (1,1) \to (1,2) \rangle + \ldots + \langle (1,n-1) \to (1,n) \rangle + \\ + \langle (1,n) \to (2,n) \rangle + \ldots + \langle (k-1,n) \to (k,n) \rangle + \langle (k,n) \to (k,n+1) \rangle,$$

see Figure 2. These relations, considered as elements of the Picard group of $\widetilde{P}$, form a basis in it. The roof relation is a pull-back to $\widetilde{P}$ of a generator of the Picard group of P. We introduce variables q_i, $i \in [1, (k-1)(n-1)]$, corresponding to box relations, and a variable q corresponding to the roof relation.



FIGURE 2. Relations

Now we describe a nef-partition corresponding to a complete intersection in the Grassmannian G. For a fixed $s \in [1, k-1]$ the *s-th horizontal basic block* is a set of all arrows $\langle (s,j) \to (s+1,j) \rangle$ with $j \in [1,n]$. Similarly, for a fixed $s \in [1, n-1]$ the *s-th vertical basic block* is a set of all arrows $\langle (i,s) \to (i,s+1) \rangle$ with $i \in [1,k]$. We also define the 0-th horizontal basic block as the set that consists of a single arrow $\langle (0,1) \to (1,1) \rangle$, and we define the n-th vertical basic block as the set that consists of a single arrow $\langle (k,n) \to (k,n+1) \rangle$.

A sum of divisors in P associated to rays corresponding to arrows in any horizontal or vertical basic block is linearly equivalent to a generator of the Picard group of P, see [BCFKS97, Proposition 4.1.4]. Thus given a complete intersection in G one can choose a nef-partition that consists of collections of rays corresponding to arrows of appropriate numbers of vertical or horizontal basic blocks.

The constant term of I-series of $\widetilde{P}$ is

$$I = I^0_{\widetilde{P}}(q, q_1, \ldots, q_{(k-1)(n-1)}).$$

In [BCFKS97, Conjecture 5.2.3] it was conjectured that

$$\widetilde{I}^G_0(q) = I(q, 1, \ldots, 1).$$

This is proved for $n = 2$ in [BCFK03, Proposition 3.5] and for any $n \geqslant 2$ in [MR13].

Consider a smooth Fano complete intersection Y in G. Let $LG_0(Y)$ be a Givental's Landau–Ginzburg model constructed for $\widetilde{P}$ and a nef-partition associated Y. Denote it's Givental's integral by I_Y^0. In discussion after Conjecture 5.2.1 in [BCFKS98] it is explained that, assuming the latter assertion, one has

$$\widetilde{I}_0^Y = I_Y^0,$$

which can be viewed as an analog of Theorem 3.8 in this particular non-toric case.

Further in Section 5 we will study the case of complete intersections in a Grassmannian of planes. For this case there is an explicit formula for constant term of regularized I-series (and thus for Givental's integral). Let

$$\gamma(r) = \sum_{i\in[1,r]} \frac{1}{i}.$$

Theorem 4.1 ([BCFK03, Proposition 3.5])**.** *Let*

$$Y = \mathrm{Gr}(2, k+2) \cap Y_1 \cap \ldots \cap Y_l$$

be a smooth Fano complete intersection with $\deg Y_i = d_i$, $\sum d_i < k+2$. *Denote*

$$d_0 = k + 2 - \sum d_i.$$

Then

$$\widetilde{I}_0^Y = \sum_{d\geqslant 0} \frac{\prod_{i=0}^l (d_0 d_i)!}{d!^{k+2}} \cdot \frac{(-1)^d}{2} \cdot \sum_{r=0}^{d} \binom{d}{r}^{k+2} \big((k+2)\,(d-2r)\,(\gamma(r) - \gamma(d-r)) - 2\big) \cdot t^{d_0 d}.$$

Summarizing, one can deal with a Grassmannian and a complete intersection therein just replacing the Grassmannian by its small toric degeneration and applying Givental's procedure to it.

Now we write down explicitly this picture after getting rid of relations as it is described in Section 3. The superpotential for G itself is the polynomial

$$a_{1,1} + \sum_{\substack{i\in[1,k-1],\\ j\in[1,n]}} \frac{a_{i+1,j}}{a_{i,j}} + \sum_{\substack{i\in[1,k],\\ j\in[1,n-1]}} \frac{a_{i,j+1}}{a_{i,j}} + \frac{1}{a_{k,n}}$$

in variables $a_{i,j}$, $i \in [1,k]$, $j \in [1,n]$, see [EHX97, B25].

Consider the following Laurent polynomials:

$$\begin{aligned} T_1 &= a_{1,1}, \\ T_{i+1} &= \sum_{j\in[1,n]} \frac{a_{i+1,j}}{a_{i,j}}, \quad i \in [1, k-1], \\ T_{k+j} &= \sum_{i\in[1,k]} \frac{a_{i,j+1}}{a_{i,j}}, \quad j \in [1, n-1], \\ T_{k+n} &= \frac{1}{a_{k,n}}. \end{aligned} \tag{4.1}$$

For any arrow

$$\alpha = \langle (i,j) \to (i',j') \rangle \in \mathrm{Ar}(\mathcal{Q}_0)$$

we define $h(\alpha)$ and $t(\alpha)$ as the vertices (i,j) and (i',j'), respectively. One can see that Laurent monomials appearing in (4.1) are of the form $a_{h(\alpha)}/a_{t(\alpha)}$ for some $\alpha \in \mathrm{Ar}(\mathcal{Q}_0)$, and Laurent polynomials listed in (4.1) are of the form

$$\sum_{\alpha\in B} \frac{a_{h(\alpha)}}{a_{t(\alpha)}},$$

where $B \subset \mathrm{Ar}(\mathcal{Q}_0)$ is some basic block.

Consider a smooth Fano complete intersection

$$Y = G \cap Y_1 \cap \ldots \cap Y_l$$

with $\deg(Y_p) = d_p$. Choose a splitting $[1, k+n] = E_0 \sqcup E_1 \sqcup \ldots \sqcup E_l$ with $|E_p| = d_p$, $p \in [1,l]$, so that $|E_0| = k+n-\sum d_p$. Define $\Sigma_p = \sum_{i\in E_p} T_i$, $p \in [0,l]$. Then the equations of anticanonical Givental's Landau–Ginzburg model for Y are

$$\Sigma_p = 1, \quad p \in [1,l], \tag{4.2}$$

and the superpotential is Σ_0.

5. Main theorem

Now we choose a specific nef-partition we are going to use in our main theorem, i. e. in Theorem 5.3 below. Informally, for any hypersurface we take a union of a suitable number of consecutive basic blocks. To make it more precise we introduce some additional terminology.

A *horizontal block* of size d is a union of d consecutive basic horizontal blocks. A *vertical block* of size d is a union of d consecutive basic vertical blocks. A *mixed block* of size d is a union of d_1 consecutive basic horizontal blocks including the $(k-1)$-th one and d_2 consecutive basic vertical blocks including the first one, where $d_1 + d_2 = d$. By a block we will mean either a horizontal block, or a vertical block, or a mixed block.



Figure 3. Blocks for the Grassmannian $\mathrm{Gr}(2,4)$

Example 5.1. Figure 3 represents several examples of blocks in a quiver corresponding to the Grassmannian $\mathrm{Gr}(2,4)$. Namely, Figures 3(a), 3(b) and 3(c) represent horizontal blocks, that are the first basic horizontal block, the 0-th basic horizontal block and a

horizontal block of size 2, respectively. Figure 3(d) represents a mixed block of size 3. Finally, Figure 3(e) represents the first basic vertical block.

The set of vertices of a block B is the set $\mathrm{Ver}(B) \subset \mathrm{Ver}(\mathcal{Q}_0)$ such that for any $v \in \mathrm{Ver}(B)$ there is an arrow $\alpha \in B$ with either $t(\alpha) = v$ or $h(\alpha) = v$. We say that an arrow $\alpha \in \mathrm{Ar}(\mathcal{Q}_0)$ is an *inner arrow* of a block B, if $t(\alpha) \in \mathrm{Ver}(B)$ and $h(\alpha) \in \mathrm{Ver}(B)$, while $\alpha \notin B$. We denote the set of inner arrows for B by $\mathrm{In}(B)$.

An *admissible quiver* $\mathcal{Q}$ is a subquiver of $\mathcal{Q}_0$ with a set of vertices $\mathrm{Ver}(\mathcal{Q}) = \mathrm{Ver}(\mathcal{Q}_0)$, and a non-empty set of arrows $\mathrm{Ar}(\mathcal{Q}) = \mathrm{Ar}(\mathcal{Q}_0) \setminus B$, where B is either a horizontal or a mixed block, and B contains the arrow $\langle (0,1) \to (1,1) \rangle$. In particular, if $\mathcal{Q}$ is an admissible quiver and $B' \subset \mathrm{Ar}(\mathcal{Q}_0)$ is a block, then

$$B'' = B' \cap \mathrm{Ar}(\mathcal{Q})$$

is again a block. Note also that if $\mathcal{Q}$ is an admissible quiver such that $\mathrm{Ar}(\mathcal{Q})$ contains the arrow $\langle (0,1) \to (1,1) \rangle$, then $\mathcal{Q} = \mathcal{Q}_0$.

Let $V = \{x_1, \ldots, x_N\}$ be a finite set. We denote the torus

$$\mathrm{Spec}\, \mathbb{C}[x_1^{\pm 1}, \ldots, x_N^{\pm 1}] \cong (\mathbb{C}^*)^N$$

by $\mathbb{T}(V)$. Note that $x_1, \ldots, x_N$ may be interpreted as coordinates on $\mathbb{T}(V)$.

A *triplet* is a collection $(\mathcal{Q}, V, R)$, where $\mathcal{Q}$ is an admissible quiver, V is a finite set of variables, R is a map from the set $\mathrm{Ver}(\mathcal{Q})$ to the set of rational functions in the variables of V.

A rational function associated to a triplet $(\mathcal{Q}, V, R)$ and a non-empty subset $C \subset \mathrm{Ar}(\mathcal{Q})$ is the rational function in the variables of V defined as

$$F_{\mathcal{Q},V,R,C} = \sum_{\alpha \in C} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}.$$

A hypersurface $H_{\mathcal{Q},V,R,C} \subset \mathbb{T}(V)$ associated to $(\mathcal{Q}, V, R)$ and C is defined by the equation $F_{\mathcal{Q},V,R,C} = 1$. A rational function associated to a triplet $(\mathcal{Q}, V, R)$ is the rational function in the variables of V defined as

$$F_{\mathcal{Q},V,R} = F_{\mathcal{Q},V,R,\mathrm{Ar}(\mathcal{Q})} = \sum_{\alpha \in \mathrm{Ar}(\mathcal{Q})} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}.$$

A change of variables that agrees with a triplet $(\mathcal{Q}, V, R)$ and with a block B is a rational map

$$\psi \colon \mathbb{T}(V') \dashrightarrow \mathbb{T}(V),$$

where V' is a set of variables such that $|V'| = |V| - 1$, the closure of the image of $\mathbb{T}(V')$ with respect to ψ is the hypersurface

$$H_{\mathcal{Q},V,R,B} \subset \mathbb{T}(V),$$

and ψ gives a birational map between $\mathbb{T}(V')$ and $H_{\mathcal{Q},V,R,B}$. By a small abuse of terminology we will sometimes omit either a triplet or a block when speaking about a change of variables that agrees with something. We will sometimes also refer to automorphisms of tori as changes of variables, but in such situations we will not mention any triplets or blocks.

Let ψ be a change of variables that agrees with a triplet $(\mathcal{Q}, V, R)$ and with a block B. *A transformation of a triplet* $(\mathcal{Q}, V, R)$ *associated to* ψ is a triplet $(\mathcal{Q}', V', R')$, where $\mathcal{Q}' \subset \mathcal{Q}$ is a quiver with $\mathrm{Ver}(\mathcal{Q}') = \mathrm{Ver}(\mathcal{Q})$ and

$$\mathrm{Ar}(\mathcal{Q}') = \mathrm{Ar}(\mathcal{Q}) \setminus B,$$

the set V' is a set of variables such that $|V'| = |V| - 1$, and $R'(i,j) = \psi^* R(i,j)$.

Remark 5.2. Let $(\mathcal{Q}, V, R)$ be a triplet, B be a block, and ψ be a change of variables that agrees with the triplet $(\mathcal{Q}, V, R)$ and with the block B. Let $(\mathcal{Q}', V', R')$ be a transformation of a triplet $(\mathcal{Q}, V, R)$ associated to ψ. Then $\psi^* F_{\mathcal{Q},V,R,B} = 1$ and

$$\psi^* F_{\mathcal{Q},V,R} = F_{\mathcal{Q}',V',R'} + 1.$$

Now we can reformulate the description of Landau–Ginzburg models for complete intersections in Grassmannians discussed in Section 4 in terms introduced above. Let

$$V_0 = \{a_{i,j}\}, \quad i \in [1,k], j \in [1,n].$$

Put $R_0(i,j) = a_{i,j}$ for $i \in [1,k]$, $j \in [1,n]$, and $R_0(0,1) = R_0(k,n+1) = 1$, see Figure 4. Let

$$Y = G \cap Y_1 \cap \ldots \cap Y_l$$

be a smooth Fano complete intersection. Let $B_1, \ldots, B_l$ be disjoint horizontal, mixed or vertical blocks such that B_i, $i \in [1,l]$, is a block of size $\deg Y_i$. Put

$$C = \mathrm{Ar}(\mathcal{Q}_0) \setminus \left(\cup_{i \in [1,l]} B_i \right).$$

Then a variety that is a complete intersection of hypersurfaces $H_{\mathcal{Q}_0,V_0,R_0,B_i}$, $i \in [1,l]$, in $\mathbb{T}(V_0)$ equipped with a function $F_{\mathcal{Q}_0,V_0,R_0,C}$ as superpotential is a Landau–Ginzburg model of Y suggested in [BCFKS98], cf. equations (4.2). Theorem 5.3 states that for given $d_1, \ldots, d_l$ there is a choice of blocks $B_1, \ldots, B_l$ and a sequence of l changes of variables such that the Landau–Ginzburg model in is fact birational to a torus, and a birational equivalence can be chosen so that the superpotential becomes a Laurent polynomial on this torus.



FIGURE 4. Triplet $(\mathcal{Q}_0, V_0, R_0)$ for the Grassmannian $\mathrm{Gr}(2,5)$

Theorem 5.3. *Let $n = 2$. Consider the triplet $(\mathcal{Q}_0, V_0, R_0)$. Let $d_1, \dots, d_l$ be positive integers such that for some $i_0 \in [0, l]$ one has $d_1, \dots, d_{i_0} > 1$ and $d_{i_0+1} = \dots = d_l = 1$. Suppose that*

$$\sum d_i < k + 2.$$

Then there exist blocks $B_1, \dots, B_l$, a sequence of triplets

$$(\mathcal{Q}_i, V_i, R_i), \quad i \in [1, l],$$

and a sequence of changes of variables

$$\psi_i \colon \mathbb{T}(V_i) \dashrightarrow \mathbb{T}(V_{i-1}), \quad i \in [1, l],$$

such that

- *the size of the block B_i is d_i;*
- *one has $B_i \cap B_j = \varnothing$ for $i \neq j$, $i, j \in [1, l]$;*
- *the change of variables ψ_i agrees with the triplet $(\mathcal{Q}_{i-1}, V_{i-1}, R_{i-1})$ and the block B_i;*
- *the triplet $(\mathcal{Q}_i, V_i, R_i)$ is a transformation of the triplet $(\mathcal{Q}_{i-1}, V_{i-1}, R_{i-1})$ associated to ψ_i;*
- *the rational function*

$$F_{\mathcal{Q}_i, V_i, R_i} = (\psi_i \circ \dots \circ \psi_1)^* F_{\mathcal{Q}_0, V_0, R_0}$$

  is a Laurent polynomial in variables of V_i.

In particular, the rational function $F_{\mathcal{Q}_l, V_l, R_l}$ is a Laurent polynomial in $2k - l$ variables.

We will prove Theorem 5.3 in Section 9. In order to do this we will write down separately the changes of variables that agree with horizontal, mixed and vertical blocks in Sections 6, 7 and 8, respectively. In most of the cases (except for a relatively easy Lemma 6.10) a change of variables will be performed in two steps. First we will choose some variable (which we will later refer to as *weight variable*), and make a monomial change of coordinates multiplying each variable by a suitably chosen power of the weight variable. After this we will exclude another variable (which we will later refer to as *main variable*) using the equation of the hypersurface associated to the triplet and the block, and check that after the corresponding substitution the Laurent polynomial associated to the triplet remains a Laurent polynomial. One effect that still looks surprising to us is that the case of a horizontal block of size 1 (i. e. of a basic horizontal block) is treated differently from the case of a horizontal block of size at least 2, so that the assertions of Lemmas 6.10 and 6.8 appear to be different indeed. Finally, since the proofs of the lemmas in Sections 6, 7 and 8 look rather messy, we illustrate them in Sections 11 and 12 by a large (and hopefully representative) sample of examples; we suspect that this may be more instructive than reading the proofs themselves.

6. Horizontal blocks

In this section we write down changes of variables that agree with horizontal blocks for Grassmannians $\mathrm{Gr}(2, k + 2)$.

We start with introducing some additional auxiliary notions.



FIGURE 5. Starting triplet for the Grassmannian $\mathrm{Gr}(2,5)$

Definition 6.1. Let V be some collection of variables. Let $W \subset V$ be a subset, and $\Lambda\colon W \to \mathbb{Z}$ be an arbitrary function. Let μ be a Laurent monomial in the variables of V. We define the Λ-*degree* of μ as

$$\deg_\Lambda(\mu) = \sum_{a \in W} \Lambda(a) \cdot \deg_a(\mu).$$

By a *total degree* of μ with respect to the variables of W we mean the Λ-degree of μ for the function $\Lambda \equiv 1$, i. e. the sum of degrees of μ with respect to the variables of W.

Definition 6.2. Let $s \in [1,k]$. Put

$$W_{\varnothing,\varnothing,s} = \{a_{i,j} \mid i \in [1,s], j \in [1,2]\}$$

if $s < k$, and put

$$W_{\varnothing,\varnothing,k} = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1,k-1]\}.$$

We define

$$\Lambda_{\varnothing,\varnothing,s}\colon W_{\varnothing,\varnothing,s} \to \mathbb{Z}$$

as $\Lambda_{\varnothing,\varnothing,s}(a_{i,2}) = i - s$ for $i \in [1,s]$ and $\Lambda_{\varnothing,\varnothing,s}(a_{i,1}) = i - s + 1$ for $i \in [1,s]$, $i \neq s-1$. Finally, we put $\Lambda_{\varnothing,\varnothing,s}(a_{s-1,1}) = 1$.

Now we start to describe our changes of variables.

Lemma 6.3. *Suppose that $n = 2$. Let $(\mathcal{Q}, V, R)$ be a triplet, and $B \subset \mathrm{Ar}(\mathcal{Q})$ be a horizontal block such that the arrow $\langle (0,1) \to (1,1) \rangle$ is contained in B. Suppose that V is a set of variables*

$$V = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1,k-1]\} \cup \{a\},$$

and the following conditions hold:

(i) *one has $R(k,2) = 1$;*
(ii) *$R(0,1) = R(k,3) = a$;*
(iii) *for $i \in [1,k]$, $j \in [1,2]$, $(i,j) \neq (k,2)$ one has $R(i,j) = a_{i,j}$, see Figure 5.*

Then there exists a change of variables ψ that agrees with the triplet $(\mathcal{Q}, V, R)$ and with the block B with the following properties. Let $(\mathcal{Q}'', V'', R'')$ be the transformation of the triplet $(\mathcal{Q}, V, R)$ associated to ψ, and let s be the largest number such that $(s, 1) \in \mathrm{Ver}(B)$. We can assume that V'' is a set of variables

$$V'' = \{a''_{i,1} \mid i \in [1, k]\} \cup \{a''_{i,2} \mid i \in [1, k-1]\}.$$

Then $\psi^* F_{\mathcal{Q},V,R}$ is a Laurent polynomial in the variables of V'' and the following assertions hold:

- (I) *the quiver $\mathcal{Q}''$ does not contain vertical arrows α such that $h(\alpha) = (i, j)$ for $i \in [1, s]$, $j \in [1, 2]$;*
- (II) *one has $R''(k, 2) = 1$;*
- (III) *for (i, j) with $i \in [s, k]$, $j \in [1, 2]$, $(i, j) \neq (k, 2)$, one has $R''(i, j) = a''_{i,j}$;*
- (IV) *for any $i \in [1, s-2]$ one has $R''(i, 1) = a''_{i,1} \cdot \bar{R}''(i)$;*
- (V) *one has $R''(s-1, 1) = \bar{R}''(i)$;*
- (VI) *for any $i \in [1, s-1]$ one has $R''(i, 2) = a''_{i,2} \cdot \bar{R}''(i)$;*
- (VII) *the rational function $R''(k, 3)$ is a Laurent polynomial in the variables of V'' such that $R''(k, 3)$ does not depend on variables $a''_{i,j}$ with $i \in [s+1, k]$, $j \in [1, 2]$, and each of its Laurent monomials has non-negative degree in each of the variables $a''_{s,j}$, $j \in [1, 2]$;*
- (VIII) *if $s < k$, then the total degree of any Laurent monomial of $R''(k, 3)$ with respect to variables $a''_{i,2}$, $i \in [1, s]$, is non-positive; if $s = k$, then the total degree of any Laurent monomial of $R''(k, 3)$ with respect to variables $a''_{i,2}$, $i \in [1, k-1]$, is non-positive;*
- (IX) *the $\Lambda_{\varnothing,\varnothing,s}$-degree of any Laurent monomial of $R''(k, 3)$ equals 1.*

Proof. If the block B consists of a single arrow $\langle (0, 1) \to (1, 1) \rangle$, then equation $F_{\mathcal{Q},V,R,B} = 1$ is equivalent to $a = a_{1,1}$. In this case we use the latter equation to exclude the variable a, and make a change of variables

$$a_{i,j} = a''_{i,j}, \quad i \in [1, k], j \in [1, 2], (i, j) \neq (k, 2).$$

Put

$$V'' = \{a''_{i,1} \mid i \in [1, k]\} \cup \{a''_{i,2} \mid i \in [1, k-1]\}.$$

We define $\psi \colon \mathbb{T}(V'') \dashrightarrow \mathbb{T}(V)$ to be the change of variables from $a_{i,j}$ to $a''_{i,j}$. We define the quiver $\mathcal{Q}''$ so that $\mathrm{Ver}(\mathcal{Q}'') = \mathrm{Ver}(\mathcal{Q})$ and $\mathrm{Ar}(\mathcal{Q}'')$ consists of all arrows of $\mathrm{Ar}(\mathcal{Q})$ except for the arrow $\langle (0, 1) \to (1, 1) \rangle$. Finally, we put $R''(i, j) = \psi^* R(i, j)$. Now the assertion of the lemma is obvious. Therefore, we assume that the size of the block B is greater than 1, so that $s \geqslant 2$.

Abusing notation a little bit, we assign $a_{k,2} = R(k, 2) = 1$; we do not mean that $a_{k,2}$ is a variable in this case (in particular, we will ignore it while computing total degrees with respect to any collection of variables), but this helps us to keep formulas more neat. Equation $F_{\mathcal{Q},V,R,B} = 1$ is equivalent to

$$\frac{a_{1,1}}{a} = 1 - \sum_{\substack{\alpha \in B, \\ \alpha \neq \langle (0,1) \to (1,1) \rangle}} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = 1 - \sum_{i \in [1, s-1],\ j \in [1,2]} \frac{a_{i+1,j}}{a_{i,j}}. \tag{6.1}$$

We choose $a_{s-1,1}$ to be the weight variable and a to be the main variable.

To start with, we make the following change of variables of V. We put $a_{s-1,1} = a'_{s-1,1}$ and we put

$$a_{i,j} = a'_{i,j} \cdot \big(a'_{s-1,1}\big)^{\mathrm{wt}(i,j)}, \quad a' = a \cdot \big(a'_{s-1,1}\big)^{\mathrm{wt}(0,1)} \tag{6.2}$$

for the following choice of weights $\mathrm{wt}(i,j)$, $(i,j) \neq (s-1,1)$. For any (i,j) with $i \in [1,s]$, $j \in [1,2]$, and for $(i,j) = (0,1)$ we put

$$\mathrm{wt}(i,j) = s - i. \tag{6.3}$$

For any (i,j) with $i \in [s+1,k]$, $j \in [1,2]$, we put $\mathrm{wt}(i,j) = 0$. In particular, this gives $\mathrm{wt}(k,2) = 0$, so that we can define $a'_{k,2} = a_{k,2} = 1$. Also, (6.3) implies that $\mathrm{wt}(s-1,1) = 1$, although we don't mean to use $\mathrm{wt}(s-1,1)$ in (6.2). Note that for any arrow $\alpha \in B$ one has

$$\mathrm{wt}\big(t(\alpha)\big) = \mathrm{wt}\big(h(\alpha)\big) + 1,$$

and for any $\alpha \in \mathrm{In}(B)$ one has $\mathrm{wt}(t(\alpha)) = \mathrm{wt}(h(\alpha))$. In particular, the weight of any non-trivial Laurent monomial appearing on the right hand side of (6.1) equals -1.

Put

$$V' = \{a'_{i,1} \mid i \in [1,k]\} \cup \{a'_{i,2} \mid i \in [1,k-1]\} \cup \{a'\}.$$

Define a collection of variables $W'_{\varnothing,\varnothing,s}$ and a function

$$\Lambda'_{\varnothing,\varnothing,s} \colon W'_{\varnothing,\varnothing,s} \to \mathbb{Z}$$

replacing the variables $a_{i,j}$ by $a'_{i,j}$ in Definition 6.2. We rewrite (6.1) as

$$\frac{a'_{1,1}}{a' \cdot a'_{s-1,1}} = 1 - \frac{1}{a'_{s-1,1}} \cdot \left(a'_{s,1} + \frac{1}{a'_{s-2,1}} + \sum_{\substack{i \in [1,s-1],\ j \in [1,2] \\ (i,j) \neq (s-1,1),(s-2,1)}} \frac{a'_{i+1,j}}{a'_{i,j}} \right) \tag{6.4}$$

if $s > 2$, and as

$$\frac{1}{a' \cdot a'_{1,1}} = 1 - \frac{1}{a'_{1,1}} \cdot \left(a'_{2,1} + \frac{a'_{2,2}}{a'_{1,2}} \right) \tag{6.5}$$

if $s = 2$. Note that the total degree with respect to variables $a'_{i,2}$, $i \in [1,s]$, of any Laurent monomial appearing on the right hand side of (6.4) and (6.5) is non-positive; actually, one can make a more precise observation: the total degree with respect to variables $a'_{i,2}$, $i \in [1,s]$, of any Laurent monomial appearing on the right hand side of (6.4) and (6.5) is zero if $s < k$ and is non-positive if $s = k$ (the latter exception appearing because we ignore $a_{k,2} = 1$ when we compute the total degree). Similarly, one can check that the $\Lambda'_{\varnothing,\varnothing,s}$-degree of any non-trivial Laurent monomial appearing on the right hand side of (6.4) and (6.5) equals 1.

Put $\delta' = a'_{1,1}$ if $s > 2$, and put $\delta' = 1$ if $s = 2$. By (6.4) and (6.5) we have

$$\frac{\delta'}{a' \cdot a'_{s-1,1}} = 1 - \frac{1}{a'_{s-1,1}} \cdot \frac{P'}{M'} = \frac{M' \cdot a'_{s-1,1} - P'}{M' \cdot a'_{s-1,1}}. \tag{6.6}$$

Here P' is a polynomial that depends only on the variables $a'_{i,j}$ with $i \in [1,s]$, $j \in [1,2]$, except for $a'_{s-1,1}$, and

$$M' = \prod_{(i,j) \in \mathcal{V}} a'_{i,j},$$

where

$$\mathcal{V} = \{(i,j) \mid i \in [1, s-1], j \in [1,2], (i,j) \neq (s-1,1)\}.$$

As above, the total degree with respect to variables $a'_{i,2}$, $i \in [1,s]$, of any Laurent monomial of the ratio P'/M', and thus of any Laurent monomial appearing on the right hand side of (6.6), is non-positive. Similarly, the $\Lambda'_{\varnothing,\varnothing,s}$-degree of any Laurent monomial of the ratio P'/M', and thus of any Laurent monomial appearing on the right hand side of (6.6), equals 1.

We rewrite (6.6) as

$$a' = \frac{M' \cdot \delta'}{M' \cdot a'_{s-1,1} - P'}. \tag{6.7}$$

Now we put

$$a''_{s-1,1} = \frac{M' \cdot a'_{s-1,1} - P'}{M'}, \tag{6.8}$$

and we put $a''_{i,j} = a'_{i,j}$ for all $i \in [1,k]$, $j \in [1,2]$, such that $(i,j) \neq (s-1,1)$ (in particular, this gives $a''_{k,2} = a'_{k,2} = 1$). Then

$$a_{s-1,1} = a'_{s-1,1} = \frac{M'' \cdot a''_{s-1,1} + P''}{M''}, \tag{6.9}$$

where P'' and M'' are obtained from P' and M' by replacing the variables $a'_{i,j}$ by the corresponding variables $a''_{i,j}$, so that M'' is the monomial

$$M'' = \prod_{(i,j)\in\mathcal{V}} a''_{i,j}.$$

Again we observe that the total degree with respect to variables $a''_{i,2}$, $i \in [1,s]$, of any Laurent monomial of the ratio P''/M'', and thus of any Laurent monomial appearing on the right hand side of (6.9), is non-positive. Similarly, we define a collection of variables $W''_{\varnothing,\varnothing,s}$ and a function

$$\Lambda''_{\varnothing,\varnothing,s} \colon W''_{\varnothing,\varnothing,s} \to \mathbb{Z}$$

replacing the variables $a_{i,j}$ by $a''_{i,j}$ in Definition 6.2, and observe that the $\Lambda''_{\varnothing,\varnothing,s}$-degree of any Laurent monomial of the ratio P''/M'', and thus of any Laurent monomial appearing on the right hand side of (6.9), equals 1.

We can rewrite (6.7) as

$$a' = \frac{\delta''}{a''_{s-1,1}}, \tag{6.10}$$

where $\delta'' = a''_{1,1}$ if $s > 2$, and $\delta'' = 1$ if $s = 2$. We see that

$$\deg_{\Lambda''_{\varnothing,\varnothing,s}}(\delta'') = 2 - s.$$

By (6.2) and (6.9) one has

$$a_{i,1} = a'_{i,1} \cdot \left(a'_{s-1,1}\right)^{s-i} = a''_{i,1} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^{s-i} \tag{6.11}$$

for any $i \in [1,s]$, $i \neq s-1$. Also, (6.2) implies that

$$a_{i,2} = a'_{i,2} \cdot \left(a'_{s-1,1}\right)^{s-i} = a''_{i,2} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^{s-i} \tag{6.12}$$

for any $i \in [1, s]$. Finally, (6.2) and (6.10) imply that

$$a = a' \cdot \left(a'_{s-1,1}\right)^s = \frac{\delta''}{a''_{s-1,1}} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^s. \tag{6.13}$$

Once again we notice that the total degree with respect to variables $a''_{i,2}$, $i \in [1, s]$, of any Laurent monomial appearing on the right hand side of (6.13) is non-positive. Also, we see that the right hand side of (6.13) does not depend on variables $a''_{i,j}$ with $i \in [s+1, k]$, $j \in [1, 2]$, and each of its Laurent monomials has non-negative degree in each of the variables $a''_{s,j}$, $j \in [1, 2]$. Similarly, we compute

$$\deg_{\Lambda''_{\varnothing,\varnothing,s}} \left(\frac{\delta''}{a''_{s-1,1}}\right) = 1 - s,$$

and see that the $\Lambda''_{\varnothing,\varnothing,s}$-degree of any Laurent monomial appearing on the right hand side of (6.13) equals 1.

Equation (6.13) allows us to exclude the variable a. Now we are going to show that after making this exclusion and changing variables $a_{i,j}$ to $a''_{i,j}$ the Laurent polynomial $F_{\mathcal{Q},V,R}$ remains a Laurent polynomial.

Let $\alpha = \langle (i, 1) \to (i, 2) \rangle$ be an inner arrow for B. Suppose that $i \neq s - 1$. Then

$$\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = \frac{a_{i,2}}{a_{i,1}} = \frac{a'_{i,2}}{a'_{i,1}} = \frac{a''_{i,2}}{a''_{i,1}}. \tag{6.14}$$

If $\alpha = \langle (s-1, 1) \to (s-1, 2) \rangle$, then

$$\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = \frac{a_{s-1,2}}{a_{s-1,1}} = \frac{a'_{s-1,2} \cdot a'_{s-1,1}}{a'_{s-1,1}} = a'_{s-1,2} = a''_{s-1,2}. \tag{6.15}$$

Put

$$V'' = \{a''_{i,1} \mid i \in [1, k]\} \cup \{a''_{i,2} \mid i \in [1, k-1]\}.$$

We define

$$\psi \colon \mathbb{T}(V'') \dashrightarrow \mathbb{T}(V)$$

to be the change of variables from $a_{i,j}$ to $a''_{i,j}$. We define the quiver $\mathcal{Q}''$ so that $\mathrm{Ver}(\mathcal{Q}'') = \mathrm{Ver}(\mathcal{Q})$ and $\mathrm{Ar}(\mathcal{Q}'') = \mathrm{Ar}(\mathcal{Q}) \setminus B$. Finally, we put $R''(i, j) = \psi^* R(i, j)$.

Denote by α_f the arrow $\langle (k, 2) \to (k, 3) \rangle$. Denote by C the set of arrows $\alpha \in \mathrm{Ar}(\mathcal{Q}) \setminus \{\alpha_f\}$ such that $h(\alpha) = (i, j)$ for some $i \in [s+1, k]$, $j \in [1, 2]$. Then the set $\mathrm{Ar}(\mathcal{Q})$ is a disjoint union of the sets B, $\mathrm{In}(B)$, C and $\{\alpha_f\}$.

One has

$$\psi^* F_{\mathcal{Q},V,R} = \psi^* \left(\sum_{\alpha \in \mathrm{Ar}(\mathcal{Q})} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}\right) =$$

$$= \psi^* \left(\sum_{\alpha \in B} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in \mathrm{In}(B)} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in C} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)}\right) =$$

$$= 1 + \psi^* \left(\sum_{\alpha \in \mathrm{In}(B)} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in C} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)}\right).$$

If $\alpha \in \mathrm{In}(B)$, then $\psi^*\left(\frac{R(h(\alpha))}{R(t(\alpha))}\right)$ is a Laurent monomial in the variables of V'' by (6.14) and (6.15).

If $\alpha \in C$, then

$$\psi^*\left(\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}\right) = \psi^*\left(\frac{a_{h(\alpha)}}{a_{t(\alpha)}}\right) = \frac{a''_{h(\alpha)}}{a''_{t(\alpha)}}$$

by conditions (ii) and (iii), because the variables $a_{i,j}$ with $i \in [s,k]$, $j \in [1,2]$, were not changed when passing from V to V' and further to V''.

Finally one can notice that

$$\psi^*\left(\frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)}\right) = \psi^* R(k,3) = \psi^* a$$

is a Laurent polynomial in the variables of V'' by (6.13).

Therefore, we see that $\psi^* F_{\mathcal{Q},V,R}$ is a Laurent polynomial in the variables of V''.

Note that assertion (I) of the lemma holds by definition of $\mathcal{Q}''$. The variables $a_{i,j} \in V$ with $i \in [s,k]$, $j \in [1,2]$, were not changed when passing from V to V' and further to V''. From this we conclude that assertions (II) and (III) of the lemma hold. Assertions (IV), (V) and (VI) hold due to equations (6.11), (6.9) and (6.12), respectively. Finally, validity of assertions (VII), (VIII) and (IX) follows from (6.13). □

Remark 6.4. In Lemma 6.3 we worked with a hypersurface given by equation

$$1 - F_{\mathcal{Q},V,R,B} = 0.$$

However, in the proof of Proposition 10.5 we will need to analyze a tubular neighborhood of the latter hypersurface, i. e. to work with an equation $1 - F_{\mathcal{Q},V,R,B} = U$. In this situation equation (6.7) takes the form

$$a' = \frac{M' \cdot \delta'}{(1-U) \cdot M' \cdot a'_{s-1,1} - P'}. \tag{6.16}$$

In the rest of this section, as well as in Sections 7 and 8, we will have to keep track of the history of changes of variables performed up to some point. This will be done with the help of the following definitions.

Definition 6.5. Let $n = 2$. Let $\mathcal{Q}$ be an admissible quiver, and let $r \in [1,k]$. We say that $(\mathcal{M}, \mathcal{W}, \gamma)$, where $\gamma \in [1,r]$ and $\mathcal{M}$ and $\mathcal{W}$ are subsets of $[1,\gamma-1]$, is a *block history* of $(\mathcal{Q},r)$ if the following conditions hold:

- one has $\mathcal{W} = \varnothing$ if and only if $\gamma = 1$ (so that one also has $\mathcal{M} = \varnothing$ in this case);
- if $\mathcal{W} \neq \varnothing$, then $|\mathcal{M}| = |\mathcal{W}| - 1$, and one has $\mathcal{M} = \{m_1, \ldots, m_{|\mathcal{M}|}\}$ and $\mathcal{W} = \{w_0, w_1, \ldots, w_{|\mathcal{M}|}\}$ with

$$w_0 < w_0 + 1 = m_1 < w_1 < \ldots < w_{|\mathcal{M}|-1} + 1 = m_{|\mathcal{M}|} < w_{|\mathcal{M}|} < w_{|\mathcal{M}|} + 1 = \gamma.$$

Definition 6.6. Let $r \in [1,k]$, and let $(\mathcal{M}, \mathcal{W}, \gamma)$ be a block history of $(\mathcal{Q},r)$. Put

$$W_{\mathcal{M},\mathcal{W},\gamma,r} = \{a_{i,1} \mid i \in [1,r]\} \cup \{a_{i,2} \mid i \in [1,\gamma-1] \setminus \mathcal{M}\} \cup \{a_{r,2}\}$$

if $r < k$, and put

$$W_{\mathcal{M},\mathcal{W},\gamma,k} = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1,\gamma-1] \setminus \mathcal{M}\}$$

if $r = k$. For any $i < \max \mathcal{W}$ (i. e. for any $i < \gamma - 1$) we put $w(i) = \min\{w \in \mathcal{W} \mid w > i\}$. We define a function

$$\Lambda_{\mathcal{M},\mathcal{W},\gamma,r} \colon W_{\mathcal{M},\mathcal{W},\gamma,r} \to \mathbb{Z}$$

as follows. If $i \in \mathcal{W}$, then we put $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{i,1}) = 1$ and $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{i,2}) = -1$. If $i \notin \mathcal{W}$ and $i < \gamma - 1$, then we put

$$\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{i,1}) = i - w(i)$$

and

$$\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{i,2}) = i - w(i) - 1.$$

If $\gamma \leqslant i \leqslant r$, then we put $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{i,1}) = 1$. Finally, if $r < k$, then we put $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{r,2}) = 0$.

If the variables of the set $W_{\mathcal{M},\mathcal{W},\gamma,r}$ are clearly labeled by some set of indices $\{(i,j)\}$ we will sometimes write $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(i,j)$ instead of $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}(a_{i,j})$.

Note that Definition 6.2 is a particular case of Definition 6.6 for $\mathcal{M} = \mathcal{W} = \varnothing$ and $\gamma = r = s$.

The following elementary observation will be rather useful for the remaining lemmas of this section.

Remark 6.7. Let V be a set of variables, and F be a Laurent polynomial in the variables of V. Let V' be some other set of variables. Consider a rational map $\psi \colon \mathbb{T}(V') \dashrightarrow \mathbb{T}(V)$. Let $W \subset V$ and $W' \subset V'$ be some subsets of variables. Choose two functions $\Lambda \colon W \to \mathbb{Z}$ and $\Lambda' \colon W' \to \mathbb{Z}$. Suppose that the rational function $\psi^* F$ is a Laurent polynomial in the variables of V'. Suppose that for any $a \in W$ the rational function $\psi^* a$ is a Laurent polynomial in the variables of V', and for any Laurent monomial μ' of $\psi^* a$ one has

$$\deg_{\Lambda'}(\mu') = \deg_{\Lambda}(a).$$

Then for any Laurent monomial μ of F one has

$$\deg_{\Lambda'}(\psi^* \mu) = \deg_{\Lambda}(\mu).$$

Now we return to our changes of variables.

Lemma 6.8. *Suppose that $n = 2$. Let $(\mathcal{Q}, V, R)$ be a triplet, and $B \subset \mathrm{Ar}(\mathcal{Q})$ be a horizontal block such that the arrow $\langle (0,1) \to (1,1) \rangle$ is not contained in B. Let r be the smallest number such that $(r,1) \in \mathrm{Ver}(B)$. Suppose that the size of the block B is greater than 1, so that B is not a basic block.*

Suppose that there is a block history $(\mathcal{M}, \mathcal{W}, r)$, i. e. one with $\gamma = r$ in the notation of Definition 6.5, of $(\mathcal{Q}, r)$ such that V is a set of variables

$$V = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1, k-1] \setminus \mathcal{M}\}.$$

Suppose that there are rational functions $\bar{R}(i)$, $i \in [1, r-1]$, in the variables of V such that the following conditions hold:

(i) *the quiver $\mathcal{Q}$ does not contain vertical arrows α such that $h(\alpha) = (i,j)$ for $i \in [1,r]$, $j \in [1,2]$;*
(ii) *one has $R(k,2) = 1$ (in what follows, we assign $a_{k,2} = R(k,2) = 1$, abusing notation a little bit);*
(iii) *for (i,j) with $i \in [r,k]$, $j \in [1,2]$, $(i,j) \neq (k,2)$, one has $R(i,j) = a_{i,j}$;*
(iv) *for any $i \in [1, r-1] \setminus \mathcal{W}$ one has $R(i,1) = a_{i,1} \cdot \bar{R}(i)$;*
(v) *for any $i \in \mathcal{W}$ one has $R(i,1) = \bar{R}(i)$;*

(vi) *for any* $i \in [1, r-1] \setminus \mathcal{M}$ *one has* $R(i,2) = a_{i,2} \cdot \bar{R}(i)$;

(vii) *for any* $i \in \mathcal{M}$ *one has*

$$R(i,2) = \frac{a_{i+1,2}}{a_{w(i),1}} \cdot \bar{R}(i),$$

where $w(i) = \min\{w \in \mathcal{W} \mid w > i\}$;

(viii) *the rational function* $R(k,3)$ *is a Laurent polynomial in the variables of* V *such that* $R(k,3)$ *does not depend on variables* $a_{i,j}$ *with* $i \in [r+1,k]$, $j \in [1,2]$, *and each of its Laurent monomials has non-negative degree in each of the variables* $a_{r,j}$, $j \in [1,2]$;

(ix) *the total degree of any Laurent monomial of* $R(k,3)$ *with respect to variables* $a_{i,2}$, $i \in [1,r] \setminus \mathcal{M}$, *is non-positive;*

(x) *the* $\Lambda_{\mathcal{M},\mathcal{W},r,r}$*-degree of any Laurent monomial of* $R(k,3)$ *equals* 1.

Then there exists a change of variables ψ *that agrees with the triplet* $(\mathcal{Q}, V, R)$ *and with the block* B *with the following properties. Let* $(\mathcal{Q}'', V'', R'')$ *be the transformation of the triplet* $(\mathcal{Q}, V, R)$ *associated to* ψ, *and let* s *be the largest number such that* $(s,1) \in \mathrm{Ver}(B)$. *Then* $\psi^* F_{\mathcal{Q},V,R}$ *is a Laurent polynomial in the variables of* V''. *Moreover, there is a block history* $(\mathcal{M}'', \mathcal{W}'', s)$ *of* $(\mathcal{Q}'', s)$ *with* $\mathcal{M}'' = \mathcal{M} \cup \{r\}$ *such that* V'' *is a set of variables*

$$V'' = \{a''_{i,1} \mid i \in [1,k]\} \cup \{a''_{i,2} \mid i \in [1,k-1] \setminus \mathcal{M}''\},$$

and conditions (i)–(x) hold after replacing $\mathcal{Q}$, V, R, $\mathcal{M}$, $\mathcal{W}$ *and* r *by* $\mathcal{Q}''$, V'', R'', $\mathcal{M}''$, $\mathcal{W}''$ *and* s, *respectively.*

Proof. Equation $F_{\mathcal{Q},V,R,B} = 1$ is equivalent to

$$\frac{a_{r+1,2}}{a_{r,2}} = 1 - \sum_{\substack{\alpha \in B, \\ \alpha \neq \langle (r,2) \to (r+1,2) \rangle}} \frac{R\big(h(\alpha)\big)}{R\big(t(\alpha)\big)} = 1 - \sum_{\substack{i \in [r,s-1],\ j \in [1,2], \\ (i,j) \neq (r,2)}} \frac{a_{i+1,j}}{a_{i,j}}. \tag{6.17}$$

We choose $a_{s-1,1}$ to be the weight variable and $a_{r,2}$ to be the main variable.

To start with, we make the following change of variables of V. We put $a_{s-1,1} = a'_{s-1,1}$ and we put

$$a_{i,j} = a'_{i,j} \cdot \big(a'_{s-1,1}\big)^{\mathrm{wt}(i,j)} \tag{6.18}$$

for the following choice of weights $\mathrm{wt}(i,j)$, $(i,j) \neq (s-1,1)$. For any (i,j) with $i \in [r,s]$, $j \in [1,2]$, we put

$$\mathrm{wt}(i,j) = s - i. \tag{6.19}$$

For any $i \in [1, r-1]$ we put $\mathrm{wt}(i,1) = 0$, and for any $i \in [1,r-1] \setminus \mathcal{M}$ we put $\mathrm{wt}(i,2) = 0$. For any (i,j) with $i \in [s+1,k]$, $j \in [1,2]$, we put $\mathrm{wt}(i,j) = 0$. In particular, this gives $\mathrm{wt}(k,2) = 0$, so that we can define $a'_{k,2} = a_{k,2} = 1$. Also (6.19) implies that $\mathrm{wt}(s-1,1) = 1$, although we don't mean to use $\mathrm{wt}(s-1,1)$ in (6.18). Note that for any arrow $\alpha \in B$ one has

$$\mathrm{wt}\big(t(\alpha)\big) = \mathrm{wt}\big(h(\alpha)\big) + 1.$$

In particular, the weight of any non-trivial Laurent monomial appearing on the right hand side of (6.17) equals -1.

Put

$$V' = \{a'_{i,1} \mid i \in [1,k]\} \cup \{a'_{i,2} \mid i \in [1,k-1] \setminus \mathcal{M}\}.$$

Put $\mathcal{M}'' = \mathcal{M} \cup \{r\}$ and $\mathcal{W}'' = \mathcal{W} \cup \{s-1\}$. Define a collection of variables $W'_{\mathcal{M}'',\mathcal{W}'',s,s}$ and a function

$$\Lambda'_{\mathcal{M}'',\mathcal{W}'',s,s} \colon W'_{\mathcal{M}'',\mathcal{W}'',s,s} \to \mathbb{Z}$$

replacing the variables $a_{i,j}$ by $a'_{i,j}$ in the appropriately applied Definition 6.6.

Note that $r < s-1$, since the size of the block B is greater than 1. We rewrite (6.17) as

$$\frac{a'_{r+1,2}}{a'_{r,2} \cdot a'_{s-1,1}} = 1 - \frac{1}{a'_{s-1,1}} \cdot \left(a'_{s,1} + \frac{1}{a'_{s-2,1}} + \sum_{\substack{i \in [r,s-1],\ j \in [1,2],\ (i,j) \neq (r,2) \\ (i,j) \neq (s-1,1),(s-2,1)}} \frac{a'_{i+1,j}}{a'_{i,j}} \right). \tag{6.20}$$

Note that the total degree with respect to variables $a'_{i,2}$, $i \in [1,s] \setminus \mathcal{M}''$, of any Laurent monomial appearing on the right hand side of (6.20) is non-positive. Similarly, one can check that the $\Lambda'_{\mathcal{M}'',\mathcal{W}'',s,s}$-degree of any non-trivial Laurent monomial appearing on the right hand side of (6.20) equals 1.

Equation (6.20) gives

$$\frac{a'_{r+1,2}}{a'_{r,2} \cdot a'_{s-1,1}} = 1 - \frac{1}{a'_{s-1,1}} \cdot \frac{P'}{M'} = \frac{M' \cdot a'_{s-1,1} - P'}{M' \cdot a'_{s-1,1}}. \tag{6.21}$$

Here P' is a polynomial that depends only on the variables $a'_{i,j}$ with $i \in [r,s]$, $j \in [1,2]$, except for $a'_{s-1,1}$ and $a'_{r,2}$, and

$$M' = \prod_{(i,j) \in \mathcal{V}} a'_{i,j},$$

where

$$\mathcal{V} = \{(i,j) \mid i \in [r,s-1], j \in [1,2], (i,j) \neq (s-1,1), (i,j) \neq (r,2)\}.$$

The total degree with respect to variables $a'_{i,2}$, $i \in [1,s] \setminus \mathcal{M}''$, of any Laurent monomial of the ratio P'/M', and thus of any Laurent monomial appearing on the right hand side of (6.21), is non-positive. Also, the $\Lambda'_{\mathcal{M}'',\mathcal{W}'',s,s}$-degree of any Laurent monomial of the ratio P'/M', and thus of any Laurent monomial appearing on the right hand side of (6.21), equals 1.

We rewrite (6.21) as

$$a'_{r,2} = \frac{M' \cdot a'_{r+1,2}}{M' \cdot a'_{s-1,1} - P'}. \tag{6.22}$$

Now we put

$$a''_{s-1,1} = \frac{M' \cdot a'_{s-1,1} - P'}{M'}, \tag{6.23}$$

we put $a''_{i,1} = a'_{i,1}$ for all $i \in [1,k]$, $i \neq s-1$, and we put $a''_{i,2} = a'_{i,2}$ for all $i \in [1,k] \setminus \mathcal{M}''$ (in particular, this gives $a''_{k,2} = a'_{k,2} = 1$). Then

$$a_{s-1,1} = a'_{s-1,1} = \frac{M'' \cdot a''_{s-1,1} + P''}{M''}, \tag{6.24}$$

where P'' and M'' are obtained from P' and M' by replacing the variables $a'_{i,j}$ by the corresponding variables $a''_{i,j}$, so that M'' is the monomial

$$M'' = \prod_{(i,j)\in\mathcal{V}} a''_{i,j}.$$

The total degree with respect to variables $a''_{i,2}$, $i \in [1,s] \setminus \mathcal{M}''$, of any Laurent monomial of the ratio P''/M'', and thus of any Laurent monomial appearing on the right hand side of (6.24), is non-positive. Define a collection of variables $W''_{\mathcal{M}'',\mathcal{W}'',s,s}$ and a function

$$\Lambda''_{\mathcal{M}'',\mathcal{W}'',s,s} \colon W''_{\mathcal{M}'',\mathcal{W}'',s,s} \to \mathbb{Z}$$

replacing the variables $a_{i,j}$ by $a''_{i,j}$ in the appropriately applied Definition 6.6. Then the $\Lambda''_{\mathcal{M}'',\mathcal{W}'',s,s}$-degree of any Laurent monomial of the ratio P''/M'', and thus of any Laurent monomial appearing on the right hand side of (6.24), equals 1. We can rewrite (6.22) as

$$a'_{r,2} = \frac{a''_{r+1,2}}{a''_{s-1,1}}. \tag{6.25}$$

Note that one has

$$a_{r,1} = a'_{r,1} \cdot \left(a'_{s-1,1}\right)^{s-r} = a''_{r,1} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^{s-r} \tag{6.26}$$

by (6.18) and (6.24). Furthermore, one has

$$a_{r,2} = a'_{r,2} \cdot \left(a'_{s-1,1}\right)^{s-r} = \frac{a''_{r+1,2}}{a''_{s-1,1}} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^{s-r} \tag{6.27}$$

by (6.18) and (6.25). Since M'' does not depend on the variables $a''_{s,j'}$, $j' \in [1,2]$, this means that for any $j \in [1,2]$ the variable $a_{r,j}$ is a Laurent polynomial in variables $a''_{i',j'}$ with $i' \in [r,s]$, $j' \in [1,2]$, $(i',j') \neq (r,2)$, such that any of its Laurent monomials has non-negative degree in variables $a''_{s,j'}$, $j' \in [1,2]$. The total degree with respect to variables $a'_{i,2}$, $i \in [1,s] \setminus \mathcal{M}''$, of any Laurent monomial appearing on the right hand side of (6.26) is non-positive. The total degree with respect to variables $a'_{i,2}$, $i \in [1,s] \setminus \mathcal{M}''$, of any Laurent monomial appearing on the right hand side of (6.27) does not exceed 1. One has

$$\deg_{\Lambda''_{\mathcal{M}'',\mathcal{W}'',s,s}}(a''_{r,1}) = r - s + 1$$

and

$$\deg_{\Lambda''_{\mathcal{M}'',\mathcal{W}'',s,s}} \left(\frac{a''_{r+1,2}}{a''_{s-1,1}}\right) = r - s,$$

so that the $\Lambda''_{\mathcal{M}'',\mathcal{W}'',s,s}$-degree of any Laurent monomial appearing on the right hand side of (6.26) equals 1, and the $\Lambda''_{\mathcal{M}'',\mathcal{W}'',s,s}$-degree of any Laurent monomial appearing on the right hand side of (6.27) equals zero. Also, (6.18) implies that

$$a_{i,1} = a'_{i,1} \cdot \left(a'_{s-1,1}\right)^{s-i} = a''_{i,1} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^{s-i} \tag{6.28}$$

for any $i \in [r,s]$, $i \neq s-1$, and

$$a_{i,2} = a'_{i,2} \cdot \left(a'_{s-1,1}\right)^{s-i} = a''_{i,2} \cdot \left(\frac{M'' \cdot a''_{s-1,1} + P''}{M''}\right)^{s-i} \tag{6.29}$$

for any $i \in [r+1, s]$.

Equation (6.27) allows us to exclude the variable $a_{r,2}$. Now we are going to show that after making this exclusion and changing variables $a_{i,j}$ to $a''_{i,j}$ the Laurent polynomial $F_{\mathcal{Q},V,R}$ remains a Laurent polynomial.

Let $\alpha = \langle (i,1) \to (i,2) \rangle$ be an inner arrow for B. Suppose that $i \neq s-1$, and $i \neq r$. Then

$$\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = \frac{a_{i,2}}{a_{i,1}} = \frac{a'_{i,2}}{a'_{i,1}} = \frac{a''_{i,2}}{a''_{i,1}}. \tag{6.30}$$

If $\alpha = \langle (s-1,1) \to (s-1,2) \rangle$, then

$$\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = \frac{a_{s-1,2}}{a_{s-1,1}} = \frac{a'_{s-1,2} \cdot a'_{s-1,1}}{a'_{s-1,1}} = a'_{s-1,2} = a''_{s-1,2}. \tag{6.31}$$

If $\alpha = \langle (r,1) \to (r,2) \rangle$, then

$$\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = \frac{a_{r,2}}{a_{r,1}} = \frac{a'_{r,2}}{a'_{r,1}} = \frac{a''_{r+1,2}}{a''_{s-1,1} \cdot a''_{r,1}}. \tag{6.32}$$

Put

$$V'' = \{a''_{i,1} \mid i \in [1,k]\} \cup \{a''_{i,2} \mid i \in [1,k-1] \setminus \mathcal{M}''\}.$$

We define

$$\psi \colon \mathbb{T}(V'') \dashrightarrow \mathbb{T}(V)$$

to be the change of variables from $a_{i,j}$ to $a''_{i,j}$. We define the quiver $\mathcal{Q}''$ so that $\mathrm{Ver}(\mathcal{Q}'') = \mathrm{Ver}(\mathcal{Q})$ and $\mathrm{Ar}(\mathcal{Q}'') = \mathrm{Ar}(\mathcal{Q}) \setminus B$. Finally, we put $R''(i,j) = \psi^* R(i,j)$.

Denote by A the set of horizontal arrows $\langle (i,1) \to (i,2) \rangle \in \mathrm{Ar}(\mathcal{Q})$ with $i \in [1, r-1]$. Denote by α_f the arrow $\langle (k,2) \to (k,3) \rangle$. Denote by C the set of arrows $\alpha \in \mathrm{Ar}(\mathcal{Q}) \setminus \{\alpha_f\}$ such that $h(\alpha) = (i,j)$ for some $i \in [s+1, k]$, $j \in [1,2]$. By condition (i) the set $\mathrm{Ar}(\mathcal{Q})$ is a disjoint union of the sets B, $\mathrm{In}(B)$, A, C and $\{\alpha_f\}$.

One has

$$\psi^* F_{\mathcal{Q},V,R} = \psi^* \left(\sum_{\alpha \in \mathrm{Ar}(\mathcal{Q})} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} \right) =$$

$$= \psi^* \left(\sum_{\alpha \in B} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in \mathrm{In}(B)} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in A} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in C} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} \right) =$$

$$= 1 + \psi^* \left(\sum_{\alpha \in \mathrm{In}(B)} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in A} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \sum_{\alpha \in C} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} \right).$$

If $\alpha \in \mathrm{In}(B)$, then $\psi^* \left(\frac{R(h(\alpha))}{R(t(\alpha))} \right)$ is a Laurent monomial in the variables of V'' by equations (6.30), (6.31) and (6.32).

If $\alpha \in A$, then $\psi^* \left(\frac{R(h(\alpha))}{R(t(\alpha))} \right)$ is a Laurent polynomial in the variables of V'' by conditions (iv)–(vii).

If $\alpha \in C$, then

$$\psi^* \left(\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} \right) = \psi^* \left(\frac{a_{h(\alpha)}}{a_{t(\alpha)}} \right) = \frac{a''_{h(\alpha)}}{a''_{t(\alpha)}}$$

by conditions (ii) and (iii), because the variables $a_{i,j}$ with $i \in [s,k]$, $j \in [1,2]$, were not changed when passing from V to V' and further to V''.

Finally one can notice that

$$\psi^* \left(\frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} \right) = \psi^* R(k,3)$$

is a Laurent polynomial in the variables of V'' by conditions (viii) and (ix), since by (6.26) and (6.27) for any $j \in [1,2]$ the rational function $\psi^* a_{r,j}$ is a Laurent polynomial in the variables $a''_{i',j'}$ with $i' \in [r,s]$, $j' \in [1,2]$, and the variables $a_{i',j'}$ with $i' \in [1, r-1] \setminus \mathcal{M}$, $j' \in [1,2]$, were not changed when passing from V to V''.

Therefore, we see that $\psi^* F_{\mathcal{Q},V,R}$ is a Laurent polynomial in the variables of V''.

Note that condition (i) of the lemma holds after replacing $\mathcal{Q}$ and r by $\mathcal{Q}''$ and s by definition of $\mathcal{Q}''$. The variables $a_{i,j} \in V$ with $i \in [s,k]$, $j \in [1,2]$, were not changed when passing from V to V' and further to V''. From this we conclude that conditions (ii) and (iii) of the lemma hold after replacing V, R and r by V'', R'' and s. Conditions (iv), (v), (vi) and (vii) hold after replacing V, R, $\mathcal{M}$, $\mathcal{W}$ and r by V'', R'', $\mathcal{M}''$, $\mathcal{W}''$ and s due to equations (6.28), (6.24), (6.29) and (6.27), respectively. Since the rational functions $\psi^* a_{r,j}$, $j \in [1,2]$, are Laurent polynomials in the variables of V'' that do not depend on the variables $a''_{i',j'}$ with $i' \in [s+1,k]$, $j' \in [1,2]$, and such that each of their Laurent monomials has non-negative degrees in the variables $a''_{s,j'}$, $j' \in [1,2]$, by (6.26) and (6.27), we see that condition (viii) of the lemma holds after replacing V, R and r by V'', R'' and s as well. Finally, by Remark 6.7 validity of conditions (ix) and (x) after replacing V, R, $\mathcal{M}$, $\mathcal{W}$ and r by V'', R'', $\mathcal{M}''$, $\mathcal{W}''$ and s also follows from equations (6.26) and (6.27). □

Remark 6.9 (cf. Remark 6.4). If in Lemma 6.8 we replace the equation $1 - F_{\mathcal{Q},V,R,B} = 0$ by $1 - F_{\mathcal{Q},V,R,B} = U$, then equation (6.22) takes the form

$$a'_{r,2} = \frac{M' \cdot a'_{r+1,2}}{(1-U) \cdot M' \cdot a'_{s-1,1} - P'}. \tag{6.33}$$

Lemma 6.10. *Suppose that $n = 2$. Let $(\mathcal{Q}, V, R)$ be a triplet, and $B \subset \mathrm{Ar}(\mathcal{Q})$ be a basic horizontal block such that the arrow $\langle (0,1) \to (1,1) \rangle$ is not contained in B. Let r be the smallest number such that $(r,1) \in \mathrm{Ver}(B)$, so that B is the r-th basic horizontal block with $r \geqslant 1$.*

Suppose that there is a block history $(\mathcal{M}, \mathcal{W}, \gamma)$ of $(\mathcal{Q}, r)$ such that V is a set of variables

$$V = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1, \gamma - 1] \setminus \mathcal{M}\} \cup \{a_{i,2} \mid i \in [r, k-1]\}.$$

Suppose that there are rational functions $\bar{R}(i)$, $i \in [1, \gamma - 1]$, in the variables of V such that the following conditions hold:

(i) *the quiver $\mathcal{Q}$ does not contain vertical arrows α such that $h(\alpha) = (i,j)$ for $i \in [1,r]$, $j \in [1,2]$;*
(ii) *one has $R(k,2) = 1$ (in what follows, we assign $a_{k,2} = R(k,2) = 1$, abusing notation a little bit);*
(iii) *for (i,j) with $i \in [r,k]$, $j \in [1,2]$, $(i,j) \neq (k,2)$, one has $R(i,j) = a_{i,j}$;*
(iv) *for any $i \in [1, \gamma - 1] \setminus \mathcal{W}$ one has $R(i,1) = a_{i,1} \cdot \bar{R}(i)$;*
(v) *for any $i \in \mathcal{W}$ one has $R(i,1) = \bar{R}(i)$;*
(vi) *for any $i \in [1, \gamma - 1] \setminus \mathcal{M}$ one has $R(i,2) = a_{i,2} \cdot \bar{R}(i)$;*

(vii) *for any* $i \in \mathcal{M}$ *one has*

$$R(i,2) = \frac{a_{i+1,2}}{a_{w(i),1}} \cdot \bar{R}(i),$$

where $w(i) = \min\{w \in \mathcal{W} \mid w > i\}$*;*

(viii) *for any* $i \in [\gamma, r-1]$ *one has*

$$R(i,1) = a_{r,1} + a_{r-1,1} + \ldots + a_{i,1};$$

(ix) *for any* $i \in [\gamma, r-1]$ *one has*

$$R(i,2) = \frac{a_{r,2} \cdot (a_{r,1} + a_{r-1,1}) \cdot (a_{r,1} + a_{r-1,1} + a_{r-2,1}) \cdot \ldots \cdot (a_{r,1} + a_{r-1,1} + \ldots + a_{i,1})}{a_{r-1,1} \cdot a_{r-2,1} \cdot \ldots \cdot a_{i,1}};$$

(x) *the rational function* $R(k,3)$ *is a Laurent polynomial in the variables of* V *such that* $R(k,3)$ *does not depend on variables* $a_{i,j}$ *with* $i \in [r+1,k]$*,* $j \in [1,2]$*, and each of its Laurent monomials has non-negative degree in each of the variables* $a_{r,j}$*,* $j \in [1,2]$*;*

(xi) *the total degree of any Laurent monomial of* $R(k,3)$ *with respect to the variables* $a_{i,2}$*,* $i \in [1,\gamma-1] \setminus \mathcal{M}$*, is non-positive;*

(xii) *the* $\Lambda_{\mathcal{M},\mathcal{W},\gamma,r}$*-degree of any Laurent monomial of* $R(k,3)$ *equals* 1*.*

Then there exists a change of variables ψ *that agrees with the triplet* $(\mathcal{Q}, V, R)$ *and with the block* B *with the following properties. Let* $(\mathcal{Q}', V', R')$ *be the transformation of the triplet* $(\mathcal{Q}, V, R)$ *associated to* ψ*. Then* $\psi^* F_{\mathcal{Q},V,R}$ *is a Laurent polynomial in the variables of* V'*. Moreover,* V' *is a set of variables*

$$V' = \{a'_{i,1} \mid i \in [1,k]\} \cup \{a'_{i,2} \mid i \in [1,\gamma-1] \setminus \mathcal{M}\} \cup \{a'_{i,2} \mid i \in [r+1, k-1]\},$$

and conditions (i)–(xii) hold after replacing $\mathcal{Q}$*,* V*,* R*, and* r *by* $\mathcal{Q}'$*,* V'*,* R'*, and* $r+1$*, respectively, and keeping* $\mathcal{M}$*,* $\mathcal{W}$ *and* γ *the same as before.*

Proof. Equation $F_{\mathcal{Q},V,R,B} = 1$ is equivalent to

$$\frac{a_{r+1,2}}{a_{r,2}} = 1 - \frac{a_{r+1,1}}{a_{r,1}},$$

which gives

$$a_{r,2} = \frac{a_{r+1,2} \cdot a_{r,1}}{a_{r,1} - a_{r+1,1}}. \tag{6.34}$$

We choose $a_{r,2}$ to be the main variable.

We make the following change of variables of V. We put

$$a'_{r,1} = a_{r,1} - a_{r+1,1},$$

we put $a'_{i,1} = a_{i,1}$ for any $i \in [1,k]$, $i \neq r$, and we put $a'_{i,2} = a_{i,2}$ for any $i \in [1,\gamma-1] \setminus \mathcal{M}$ and for any $i \in [r,k]$ (in particular, this gives $a'_{k,2} = a_{k,2} = 1$). Now one has

$$a_{r,1} = a'_{r,1} + a'_{r+1,1}, \tag{6.35}$$

and we rewrite (6.34) as

$$a_{r,2} = a'_{r,2} = \frac{a'_{r+1,2} \cdot (a'_{r,1} + a'_{r+1,1})}{a'_{r,1}}. \tag{6.36}$$

Define a collection of variables $W'_{\mathcal{M},\mathcal{W},\gamma,r+1}$ and a function

$$\Lambda'_{\mathcal{M},\mathcal{W},\gamma,r+1}\colon W'_{\mathcal{M},\mathcal{W},\gamma,r+1}\to\mathbb{Z}$$

replacing the variables $a_{i,j}$ by $a'_{i,j}$ in the appropriately applied Definition 6.6. It is straightforward to check that the $\Lambda'_{\mathcal{M},\mathcal{W},\gamma,r+1}$-degree of any Laurent monomial appearing on the right hand side of (6.35) equals 1, while the $\Lambda'_{\mathcal{M},\mathcal{W},\gamma,r+1}$-degree of any Laurent monomial appearing on the right hand side of (6.36) equals zero.

Equation (6.36) allows us to exclude the variable $a_{r,2}$. Now we are going to show that after making this exclusion and changing variables $a_{i,j}$ to $a'_{i,j}$ the Laurent polynomial $F_{\mathcal{Q},V,R}$ remains a Laurent polynomial.

Put

$$V'=\{a_{i,1}\mid i\in[1,k]\}\cup\{a_{i,2}\mid i\in[r+1,k-1]\}.$$

We define

$$\psi\colon\mathbb{T}(V')\dashrightarrow\mathbb{T}(V)$$

to be the change of variables from $a_{i,j}$ to $a'_{i,j}$. We define the quiver $\mathcal{Q}'$ so that $\mathrm{Ver}(\mathcal{Q}')=\mathrm{Ver}(\mathcal{Q})$ and $\mathrm{Ar}(\mathcal{Q}')=\mathrm{Ar}(\mathcal{Q})\setminus B$. Finally, we put $R'(i,j)=\psi^*R(i,j)$.

By construction we see that conditions (i)–(iii) of the lemma hold after replacing $\mathcal{Q}$, V, R and r by $\mathcal{Q}'$, V', R' and $r+1$, respectively. Since the variables $a_{i,1}$, $i\in[1,\gamma-1]$, and $a_{i,2}$, $i\in[1,\gamma-1]\setminus\mathcal{M}$, were not changed when passing from V to V', we see that conditions (iv)–(vii) of the lemma hold after replacing $\mathcal{Q}$, V, R and r by $\mathcal{Q}'$, V', R' and $r+1$, respectively, and keeping $\mathcal{M}$, $\mathcal{W}$ and γ the same as before. Plugging in equations (6.35) and (6.36) into conditions (viii)–(xii) of the lemma and keeping in mind Remark 6.7, we see that these conditions hold after replacing $\mathcal{Q}$, V, R and r by $\mathcal{Q}'$, V', R' and $r+1$ as well.

Denote by A_1 the set of horizontal arrows $\langle(i,1)\to(i,2)\rangle\in\mathrm{Ar}(\mathcal{Q})$ with $i\in[1,\gamma-1]$, and by A_2 the set of horizontal arrows $\langle(i,1)\to(i,2)\rangle\in\mathrm{Ar}(\mathcal{Q})$ with $i\in[\gamma,r]$. Denote by α_f the arrow $\langle(k,2)\to(k,3)\rangle$. Denote by C the set of arrows $\alpha\in\mathrm{Ar}(\mathcal{Q})\setminus\{\alpha_f\}$ such that $t(\alpha)=(i,j)$ for some $i\in[r+1,k]$, $j\in[1,2]$. By condition (i) the set $\mathrm{Ar}(\mathcal{Q})$ is a disjoint union of the sets B, A, C and $\{\alpha_f\}$.

One has

$$\psi^*F_{\mathcal{Q},V,R}=\psi^*\left(\sum_{\alpha\in\mathrm{Ar}(\mathcal{Q})}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}\right)= \tag{6.37}$$

$$=\psi^*\left(\sum_{\alpha\in B}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}+\sum_{\alpha\in A_1}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}+\sum_{\alpha\in A_2}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}+\sum_{\alpha\in C}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}+\frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)}\right)=$$

$$=1+\sum_{i\in[\gamma,r]}\frac{a'_{r,2}\cdot(a'_{r,1}+a'_{r-1,1})\cdot\ldots\cdot(a'_{r,1}+\ldots+a'_{i+1,1})}{a'_{r-1,1}\cdot a'_{r-2,1}\cdot\ldots\cdot a'_{i,1}}+$$

$$+\sum_{i\in[r+1,k]}\frac{a'_{i,2}}{a'_{i,1}}+\sum_{i\in[r+1,k-1]}\left(\frac{a'_{i+1,1}}{a'_{i,1}}+\frac{a'_{i+1,2}}{a'_{i,2}}\right)+\psi^*\left(\sum_{\alpha\in A_1}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}\right)+\psi^*R(k,3).$$

By conditions (iv)–(vii) of the lemma we know that the rational function

$$\psi^*\left(\sum_{\alpha\in A_1}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}\right)$$

is a Laurent polynomial in the variables of V'. Also, we already know that the rational function $\psi^*R(k,3)=R'(k,3)$ is a Laurent polynomial. Therefore, equation (6.37) shows that $\psi^*F_{\mathcal{Q},V,R}$ is a Laurent polynomial in the variables of V', which completes the proof of the lemma. □

Remark 6.11 (cf. Remark 6.4). If in Lemma 6.10 we replace the equation $1-F_{\mathcal{Q},V,R,B}=0$ by $1-F_{\mathcal{Q},V,R,B}=U$, then equation 6.34 takes the form

$$a_{r,2}=\frac{a_{r+1,2}\cdot a_{r,1}}{(1-U)\cdot a_{r,1}-a_{r+1,1}}. \tag{6.38}$$

7. Mixed blocks

In this section we write down changes of variables that agree with mixed blocks for Grassmannians $\mathrm{Gr}(2,k+2)$.

Lemma 7.1. *Suppose that $n=2$. Let $(\mathcal{Q},V,R)$ be a triplet, and $B\subset \mathrm{Ar}(\mathcal{Q})$ be a mixed block such that the arrow $\langle(0,1)\to(1,1)\rangle$ is contained in B and the arrow $\langle(k,2)\to(k,3)\rangle$ is not contained in B. Suppose that V is a set of variables*

$$V=\{a_{i,1}\mid i\in[1,k]\}\cup\{a_{i,2}\mid i\in[1,k-1]\}\cup\{a\},$$

and the following conditions hold:

(i) *one has $R(k,2)=1$;*
(ii) *$R(0,1)=R(k,3)=a$;*
(iii) *for $i\in[1,k]$, $j\in[1,2]$, $(i,j)\neq(k,2)$ one has $R(i,j)=a_{i,j}$, see Figure 5*

Then there exists a change of variables ψ that agrees with the triplet $(\mathcal{Q},V,R)$ and with the block B such that for the transformation $(\mathcal{Q}'',V'',R'')$ of the triplet $(\mathcal{Q},V,R)$ associated to ψ the rational function $\psi^*F_{\mathcal{Q},V,R}$ is a Laurent polynomial in the variables of V''.

Proof. Abusing notation a little bit, we assign $a_{k,2}=R(k,2)=1$. Equation $F_{\mathcal{Q},V,R,B}=1$ is equivalent to

$$\frac{a_{1,1}}{a}=1-\sum_{\substack{\alpha\in B,\\ \alpha\neq\langle(0,1)\to(1,1)\rangle}}\frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)}=1-\left(\sum_{i\in[1,k-1],\ j\in[1,2]}\frac{a_{i+1,j}}{a_{i,j}}+\sum_{i\in[1,k]}\frac{a_{i,2}}{a_{i,1}}\right). \tag{7.1}$$

We choose $a_{k-1,2}$ to be the weight variable and a to be the main variable.

To start with, we make the following change of variables of V. We put $a_{k-1,2}=a'_{k-1,2}$ and we put

$$a_{i,j}=a'_{i,j}\cdot\left(a'_{k-1,2}\right)^{\mathrm{wt}(i,j)},\quad a'=a\cdot\left(a'_{k-1,2}\right)^{\mathrm{wt}(0,1)} \tag{7.2}$$

for the following choice of weights $\mathrm{wt}(i,j)$, $(i,j)\neq(k-1,2)$. For any (i,j) with $i\in[1,k]$, $j\in[1,2]$, and for $(i,j)=(0,1)$ we put

$$\mathrm{wt}(i,j)=k+2-i-j. \tag{7.3}$$

In particular, this gives $\mathrm{wt}(k,2)=0$, so that we can define $a'_{k,2}=a_{k,2}=1$. Also, (7.3) implies that $\mathrm{wt}(k-1,2)=1$, although we don't mean to use $\mathrm{wt}(k-1,2)$ in (7.2). Note that for any arrow $\alpha\in B$ one has

$$\mathrm{wt}\big(t(\alpha)\big)=\mathrm{wt}\big(h(\alpha)\big)+1.$$

In particular, the weight of any non-trivial Laurent monomial appearing on the right hand side of (7.1) equals -1.

Put

$$V'=\{a'_{i,1}\mid i\in[1,k]\}\cup\{a'_{i,2}\mid i\in[1,k-1]\}\cup\{a'\}.$$

We rewrite (7.1) as

$$\frac{a'_{1,1}}{a'\cdot a'_{k-1,2}}=1-\frac{1}{a'_{k-1,2}}\cdot\left(\Delta'+\sum_{\substack{i\in[1,k-1],\ j\in[1,2],\\ (i,j)\neq(k-1,2),(k-2,2)}}\frac{a'_{i+1,j}}{a'_{i,j}}+\sum_{i\in[1,k]}\frac{a'_{i,2}}{a'_{i,1}}\right),\tag{7.4}$$

where

$$\Delta'=1+\frac{1}{a'_{k-2,2}}$$

if $k>2$, and $\Delta'=1$ if $k=2$. This gives

$$\frac{a'_{1,1}}{a'\cdot a'_{k-1,2}}=1-\frac{1}{a'_{k-1,2}}\cdot\frac{P'}{M'}=\frac{M'\cdot a'_{k-1,2}-P'}{M'\cdot a'_{k-1,2}}.\tag{7.5}$$

Here P' is a polynomial that does not depend on the variables a' and $a'_{k-1,2}$, and

$$M'=\prod_{(i,j)\in\mathcal{V}}a'_{i,j},$$

where

$$\mathcal{V}=\{(i,j)\mid i\in[1,k-1],j\in[1,2],(i,j)\neq(k-1,2)\}.$$

We rewrite (7.5) as

$$a'=\frac{M'\cdot a'_{1,1}}{M'\cdot a'_{k-1,2}-P'}.\tag{7.6}$$

Now we put

$$a''_{k-1,2}=\frac{M'\cdot a'_{k-1,2}-P'}{M'},\tag{7.7}$$

and we put $a''_{i,j}=a'_{i,j}$ for all $i\in[1,k]$, $j\in[1,2]$, such that $(i,j)\neq(k-1,2)$ (in particular, this gives $a''_{k,2}=a'_{k,2}=1$). Then

$$a_{k-1,2}=a'_{k-1,2}=\frac{M''\cdot a''_{k-1,2}+P''}{M''},\tag{7.8}$$

where P'' and M'' are obtained from P' and M' by replacing the variables $a'_{i,j}$ by the corresponding variables $a''_{i,j}$, so that M'' is the monomial

$$M''=\prod_{(i,j)\in\mathcal{V}}a''_{i,j}.$$

We can rewrite (7.6) as

$$a' = \frac{a''_{1,1}}{a''_{k-1,2}}. \tag{7.9}$$

Equations (7.2) and (7.9) imply that

$$a = a' \cdot \left(a'_{k-1,2}\right)^{k+1} = \frac{a''_{1,1}}{a''_{k-1,2}} \cdot \left(\frac{M'' \cdot a''_{k-1,2} + P''}{M''}\right)^{k+1}. \tag{7.10}$$

Equation (7.10) allows us to exclude the variable a. Put

$$V'' = \{a''_{i,1} \mid i \in [1,k]\} \cup \{a''_{i,2} \mid i \in [1,k-1]\}.$$

We define

$$\psi \colon \mathbb{T}(V'') \dashrightarrow \mathbb{T}(V)$$

to be the change of variables from $a_{i,j}$ to $a''_{i,j}$. We define the quiver $\mathcal{Q}''$ so that $\mathrm{Ver}(\mathcal{Q}'') = \mathrm{Ver}(\mathcal{Q})$, and the set $\mathrm{Ar}(\mathcal{Q}'')$ consists of a single arrow

$$\alpha_f = \langle (k,2) \to (k,3) \rangle.$$

Finally, we put $R''(i,j) = \psi^* R(i,j)$.

One has

$$\psi^* F_{\mathcal{Q},V,R} = \psi^* \left(\sum_{\alpha \in \mathrm{Ar}(\mathcal{Q})} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} \right) =$$

$$= \psi^* \left(\sum_{\alpha \in B} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} \right) =$$

$$= 1 + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} = 1 + \psi^* R(k,3) = 1 + \psi^* a.$$

The latter is a Laurent polynomial in the variables of V'' by (7.10). □

Remark 7.2 (cf. Remark 6.4). If in Lemma 7.1 we replace the equation $1 - F_{\mathcal{Q},V,R,B} = 0$ by $1 - F_{\mathcal{Q},V,R,B} = U$, then equation 7.6 takes the form

$$a' = \frac{M' \cdot a'_{1,1}}{(1-U) \cdot M' \cdot a'_{k-1,2} - P'} \tag{7.11}$$

Lemma 7.3. *Suppose that $n = 2$. Let $(\mathcal{Q}, V, R)$ be a triplet, and $B \subset \mathrm{Ar}(\mathcal{Q})$ be a mixed block such that the arrow $\langle (0,1) \to (1,1) \rangle$ is not contained in B. Let r be the smallest number such that there is a vertical arrow $\langle (r,1) \to (r+1,1) \rangle \in B$.*

Suppose that there is a block history $(\mathcal{M}, \mathcal{W}, r)$, i. e. one with $\gamma = r$ in the notation of Definition 6.5, of $(\mathcal{Q}, r)$ such that V is a set of variables

$$V = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1,k-1] \setminus \mathcal{M}\}.$$

Suppose that there are rational functions $\bar{R}(i)$, $i \in [1, r-1]$, in the variables of V such that the following conditions hold:

(i) *the quiver $\mathcal{Q}$ does not contain vertical arrows α such that $h(\alpha) = (i,j)$ for $i \in [1,r]$, $j \in [1,2]$;*
(ii) *one has $R(k,2) = 1$;*

(iii) *for* (i,j) *with* $i \in [r,k]$, $j \in [1,2]$, $(i,j) \neq (k,2)$, *one has* $R(i,j) = a_{i,j}$;
(iv) *for any* $i \in [1, r-1] \setminus \mathcal{W}$ *one has* $R(i,1) = a_{i,1} \cdot \bar{R}(i)$;
(v) *for any* $i \in \mathcal{W}$ *one has* $R(i,1) = \bar{R}(i)$;
(vi) *for any* $i \in [1, r-1] \setminus \mathcal{M}$ *one has* $R(i,2) = a_{i,2} \cdot \bar{R}(i)$;
(vii) *for any* $i \in \mathcal{M}$ *one has*

$$R(i,2) = \frac{a_{i+1,2}}{a_{w(i),1}} \cdot \bar{R}(i),$$

where $w(i) = \min\{w \in \mathcal{W} \mid w > i\}$;
(viii) *the rational function* $R(k,3)$ *is a Laurent polynomial in the variables of* V *such that* $R(k,3)$ *does not depend on variables* $a_{i,j}$ *with* $i \in [r+1,k]$, $j \in [1,2]$, *and each of its Laurent monomials has non-negative degree in each of the variables* $a_{r,j}$, $j \in [1,2]$;
(ix) *the total degree of any Laurent monomial of* $R(k,3)$ *with respect to variables* $a_{i,2}$, $i \in [1,r] \setminus \mathcal{M}$, *is non-positive.*

Then there exists a change of variables ψ *that agrees with the triplet* $(\mathcal{Q}, V, R)$ *and with the block* B *such that for the transformation* $(\mathcal{Q}'', V'', R'')$ *of the triplet* $(\mathcal{Q}, V, R)$ *associated to* ψ *the rational function* $\psi^* F_{\mathcal{Q},V,R}$ *is a Laurent polynomial in the variables of* V''.

Proof. Abusing notation a little bit, we assign $a_{k,2} = R(k,2) = 1$. By conditions (iv)–(vii) equation $F_{\mathcal{Q},V,R,B} = 1$ is equivalent to

$$\frac{a_{r+1,1} + a_{r,2}}{a_{r,1}} = 1 - \sum_{\substack{\alpha \in B,\ \alpha \neq \langle (r,1) \to (r,2) \rangle \\ \alpha \neq \langle (r,1) \to (r+1,1) \rangle}} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} = \tag{7.12}$$

$$= 1 - \left(\sum_{\substack{i \in [r+1,k-1] \\ j \in [1,2]}} \frac{a_{i+1,j}}{a_{i,j}} + \sum_{i \in [r+1,k]} \frac{a_{i,2}}{a_{i,1}} + \sum_{i \in [1,r-1] \setminus (\mathcal{M} \cup \mathcal{W})} \frac{a_{i,2}}{a_{i,1}} + \sum_{i \in \mathcal{W}} a_{i,2} + \sum_{i \in \mathcal{M}} \frac{a_{i+1,2}}{a_{w(i),1} \cdot a_{i,1}} \right).$$

We choose $a_{k,1}$ to be the weight variable and $a_{r,1}$ to be the main variable.

To start with, we make the following change of variables of V. We put $a_{k,1} = a'_{k,1}$ and we put

$$a_{i,j} = a'_{i,j} \cdot \left(a'_{k,1}\right)^{\mathrm{wt}(i,j)} \tag{7.13}$$

for the following choice of weights $\mathrm{wt}(i,j)$, $(i,j) \neq (k,1)$. For any (i,j) with $i \in [r,k]$, $j \in [1,2]$, we put

$$\mathrm{wt}(i,j) = k + 2 - i - j. \tag{7.14}$$

In particular, this gives $\mathrm{wt}(k,2) = 0$, so that we can define $a'_{k,2} = a_{k,2} = 1$. For any $i \in [1, r-1]$ we put $\mathrm{wt}(i,1) = 0$, and for any $i \in [1, r-1] \setminus \mathcal{M}$ we put $\mathrm{wt}(i,2) = -1$. Also, (7.14) implies that $\mathrm{wt}(k,1) = 1$, although we don't mean to use $\mathrm{wt}(k,1)$ in (7.13). Note that the weight of any non-trivial Laurent monomial appearing on the right hand side of (7.12) equals -1.

Put

$$V' = \{a'_{i,1} \mid i \in [1,k]\} \cup \{a'_{i,2} \mid i \in [1,k-1] \setminus \mathcal{M}\}.$$

We rewrite (7.12) as

$$\frac{\delta' + a'_{r,2}}{a'_{r,1} \cdot a'_{k,1}} = \tag{7.15}$$

$$= 1 - \frac{1}{a'_{k,1}} \cdot \left(\Delta' + \sum_{\substack{i \in [r+1, k-1],\ j \in [1,2], \\ (i,j) \neq (r,1),\ (i,j) \neq (k-1,1)}} \frac{a'_{i+1,j}}{a'_{i,j}} + \sum_{i \in [r+1, k-1],\ j \in [1,2]} \frac{a'_{i,2}}{a'_{i,1}} + \right.$$

$$\left. + \sum_{i \in [1, r-1] \setminus (\mathcal{M} \cup \mathcal{W})} \frac{a'_{i,2}}{a'_{i,1}} + \sum_{i \in \mathcal{W}} a'_{i,2} + \sum_{i \in \mathcal{M}} \frac{a'_{i+1,2}}{a'_{w(i),1} \cdot a'_{i,1}} \right),$$

where

$$\Delta' = 1 + \frac{1}{a'_{k-1,1}}$$

and $\delta' = a'_{r+1,1}$ if $r < k-1$, while $\Delta' = \delta' = 1$ if $r = k-1$.

Equation (7.15) gives

$$\frac{\delta' + a'_{r,2}}{a'_{r,1} \cdot a'_{k,1}} = 1 - \frac{1}{a'_{k,1}} \cdot \frac{P'}{M'} = \frac{M' \cdot a'_{k,1} - P'}{M' \cdot a'_{k,1}}. \tag{7.16}$$

Here P' is a polynomial that does not depend on the variables $a'_{k,1}$ and $a'_{r,1}$, and M' is a monomial that does not depend on the variables $a'_{k,1}$ and $a'_{r,1}$. We rewrite (7.16) as

$$a'_{r,1} = \frac{M' \cdot (\delta' + a'_{r,2})}{M' \cdot a'_{k,1} - P'}. \tag{7.17}$$

Now we put

$$a''_{k,1} = \frac{M' \cdot a'_{k,1} - P'}{M'}, \tag{7.18}$$

we put $a''_{i,1} = a'_{i,1}$ for all $i \in [1, k-1]$, $i \neq r$, and we put $a''_{i,2} = a'_{i,2}$ for all $i \in [1,k] \setminus \mathcal{M}$ (in particular, this gives $a''_{k,2} = a'_{k,2} = 1$). Then

$$a_{k,1} = a'_{k,1} = \frac{M'' \cdot a''_{k,1} + P''}{M''}, \tag{7.19}$$

where P'' and M'' are obtained from P' and M' by replacing the variables $a'_{i,j}$ by the corresponding variables $a''_{i,j}$.

We can rewrite (7.17) as

$$a'_{r,1} = \frac{\delta'' + a''_{r,2}}{a''_{k,1}}, \tag{7.20}$$

where $\delta'' = a''_{r+1,1}$ if $r < k-1$, while $\delta' = 1$ if $r = k-1$. Equations (7.13) and (7.20) imply that

$$a_{r,1} = a_{r,1} \cdot \left(a'_{k,1}\right)^{k-r+1} = \frac{\delta''}{a''_{k,1}} \cdot \left(\frac{M'' \cdot a''_{k,1} + P''}{M''} \right)^{k-r+1}. \tag{7.21}$$

Also, (7.13) implies that

$$a_{i,1} = a'_{i,1} = a''_{i,1} \tag{7.22}$$

for $i \in [1, r-1]$, while

$$a_{i,2} = a'_{i,2} \cdot \left(a'_{k,1}\right)^{-1} = a''_{i,2} \cdot \left(\frac{M'' \cdot a''_{k,1} + P''}{M''}\right)^{-1} \tag{7.23}$$

for $i \in [1, r-1] \setminus \mathcal{M}$, and

$$a_{r,2} = a'_{r,2} \cdot \left(a'_{k,1}\right)^{k-r} = a''_{r,2} \cdot \left(\frac{M'' \cdot a''_{k,1} + P''}{M''}\right)^{k-r}. \tag{7.24}$$

Equation (7.21) allows us to exclude the variable $a_{r,1}$. Put

$$V'' = \{a''_{i,1} \mid i \in [1,k], i \neq r\} \cup \{a''_{i,2} \mid i \in [1, k-1] \setminus \mathcal{M}\}.$$

We define

$$\psi \colon \mathbb{T}(V'') \dashrightarrow \mathbb{T}(V)$$

to be the change of variables from $a_{i,j}$ to $a''_{i,j}$. We define the quiver $\mathcal{Q}''$ so that $\mathrm{Ver}(\mathcal{Q}'') = \mathrm{Ver}(\mathcal{Q})$, and the set $\mathrm{Ar}(\mathcal{Q}'')$ consists of a single arrow

$$\alpha_f = \langle (k,2) \to (k,3) \rangle.$$

Finally, we put $R''(i,j) = \psi^* R(i,j)$.

One has

$$\psi^* F_{\mathcal{Q},V,R} = \psi^* \left(\sum_{\alpha \in \mathrm{Ar}(\mathcal{Q})} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} \right) =$$

$$= \psi^* \left(\sum_{\alpha \in B} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} \right) =$$

$$= 1 + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} = 1 + \psi^* R(k,3).$$

Conditions (viii) and (ix) imply that $R(k,3)$ is a Laurent polynomial in the variables of V independent of variables $a_{i,j}$ with $i \in [r+1, k]$, $j \in [1,2]$ (in particular, independent of the variable $a_{k,1}$), such that each of its Laurent monomials has a non-negative degree in the variables $a_{r,j}$, $j \in [1,2]$, and non-positive total degree with respect to variables $a_{i,2}$ with $i \in [1, r-1]$. Recall that one has $\mathrm{wt}(i,1) = 0$ for $i \in [1, r-1]$ and $\mathrm{wt}(r,1) > 0$, while $\mathrm{wt}(i,2) < 0$ for $i \in [1, r-1] \setminus \mathcal{M}$ and $\mathrm{wt}(r,2) > 0$. This means that the weight of any Laurent monomial of $R(k,3)$ is non-negative. Therefore, plugging (7.21), (7.22), (7.23) and (7.24) into $R(k,3)$, we see that $\psi^* R(k,3)$, and thus also $\psi^* F_{\mathcal{Q},V,R}$, is a Laurent polynomials in the variables of V''. □

Remark 7.4 (cf. Remark 6.4). If in Lemma 7.3 we replace the equation $1 - F_{\mathcal{Q},V,R,B} = 0$ by $1 - F_{\mathcal{Q},V,R,B} = U$, then equation 7.17 takes the form

$$a'_{r,1} = \frac{M' \cdot (\delta' + a'_{r,2})}{(1-U) \cdot M' \cdot a'_{k,1} - P'}. \tag{7.25}$$

8. Vertical blocks

In this section we write down changes of variables that agree with vertical blocks for Grassmannians $\mathrm{Gr}(2, k+2)$ and make some concluding remarks on the changes of variables that agree with various kinds of blocks.

Lemma 8.1. *Suppose that $n = 2$. Let $(\mathcal{Q}, V, R)$ be a triplet, and $B \subset \mathrm{Ar}(\mathcal{Q})$ be a vertical block such that the arrow $\langle (k,2) \to (k,3) \rangle$ is not contained in B (i. e. B is the first basic vertical block).*

Suppose that there is a block history $(\mathcal{M}, \mathcal{W}, \gamma)$ of $(\mathcal{Q}, k)$ such that V is a set of variables

$$V = \{a_{i,1} \mid i \in [1,k]\} \cup \{a_{i,2} \mid i \in [1, \gamma - 1] \setminus \mathcal{M}\}.$$

Suppose that there are rational functions $\bar{R}(i)$, $i \in [1, \gamma - 1]$, in the variables of V such that the following conditions hold:

(i) *the quiver $\mathcal{Q}$ does not contain vertical arrows;*
(ii) *one has $R(k,2) = 1$;*
(iii) *one has $R(k,1) = a_{k,1}$;*
(iv) *for any $i \in [1, \gamma - 1] \setminus \mathcal{W}$ one has $R(i,1) = a_{i,1} \cdot \bar{R}(i)$;*
(v) *for any $i \in \mathcal{W}$ one has $R(i,1) = \bar{R}(i)$;*
(vi) *for any $i \in [1, \gamma - 1] \setminus \mathcal{M}$ one has $R(i,2) = a_{i,2} \cdot \bar{R}(i)$;*
(vii) *for any $i \in \mathcal{M}$ one has*

$$R(i,2) = \frac{a_{i+1,2}}{a_{w(i),1}} \cdot \bar{R}(i),$$

where $w(i) = \min\{w \in \mathcal{W} \mid w > i\}$;
(viii) *for any $i \in [\gamma, k-1]$ one has*

$$R(i,1) = a_{k,1} + a_{k-1,1} + \ldots + a_{i,1};$$

(ix) *for any $i \in [\gamma, k-1]$ one has*

$$R(i,2) = \frac{(a_{k,1} + a_{k-1,1}) \cdot (a_{k,1} + a_{k-1,1} + a_{k-2,1}) \cdot \ldots \cdot (a_{k,1} + a_{k-1,1} + \ldots + a_{i,1})}{a_{k-1,1} \cdot a_{k-2,1} \cdot \ldots \cdot a_{i,1}};$$

(x) *the rational function $R(k,3)$ is a Laurent polynomial in the variables of V such that the $\Lambda_{\mathcal{M},\mathcal{W},\gamma,k}$-degree of any Laurent monomial of $R(k,3)$ is non-negative.*

Then there exists a change of variables ψ that agrees with the triplet $(\mathcal{Q}, V, R)$ and with the block B such that for the transformation $(\mathcal{Q}'', V'', R'')$ of the triplet $(\mathcal{Q}, V, R)$ associated to ψ the rational function $\psi^* F_{\mathcal{Q},V,R}$ is a Laurent polynomial in the variables of V''.

Proof. Abusing notation a little bit, we assign $a_{k,2} = R(k,2) = 1$. Note that conditions (iii) and (iv) imply that for any $i \in [\gamma, k]$ one has

$$\frac{R(i,2)}{R(i,1)} = \frac{(a_{k,1} + a_{k-1,1}) \cdot \ldots \cdot (a_{k,1} + a_{k-1,1} + \ldots + a_{i+1,1})}{a_{k-1,1} \cdot a_{k-2,1} \cdot \ldots \cdot a_{i+1,1}} \cdot \frac{1}{a_{i,1}}.$$

Therefore, by conditions (iv)–(ix) equation $F_{\mathcal{Q},V,R,B} = 1$ is equivalent to

(8.1)
$$\frac{(a_{k,1}+a_{k-1,1})\cdot\ldots\cdot(a_{k,1}+a_{k-1,1}+\ldots+a_{\gamma+1,1})}{a_{k-1,1}\cdot\ldots\cdot a_{\gamma+1,1}}\cdot\frac{1}{a_{\gamma,1}} = 1 - \sum_{\substack{\alpha\in B,\\ \alpha\neq\langle(k,1)\to(k,2)\rangle}} \frac{R\,(h(\alpha))}{R\,(t(\alpha))} =$$

$$= 1 - \left(\sum_{i\in[1,\gamma-1]\setminus(\mathcal{M}\cup\mathcal{W})} \frac{a_{i,2}}{a_{i,1}} + \sum_{i\in\mathcal{W}} a_{i,2} + \sum_{i\in\mathcal{M}} \frac{a_{i+1,2}}{a_{w(i),1}\cdot a_{i,1}} + \right.$$

$$\left. + \sum_{i\in[\gamma+1,k]} \frac{(a_{k,1}+a_{k-1,1})\cdot\ldots\cdot(a_{k,1}+a_{k-1,1}+\ldots+a_{i+1,1})}{a_{k-1,1}\cdot a_{k-2,1}\cdot\ldots\cdot a_{i+1,1}}\cdot\frac{1}{a_{i,1}} \right).$$

If $\gamma < k$, we put $u = k$. If $\gamma = k$, then $\mathcal{W} \neq \varnothing$, and we put $u = \min\mathcal{W}$ (so that $u \neq \gamma$ by definition of block history). In both cases we choose $a_{u,1}$ to be the weight variable and $a_{\gamma,1}$ to be the main variable.

To start with, we make the following change of variables of V. We put $a_{u,1} = a'_{u,1}$ and we put

$$a_{i,j} = a'_{i,j}\cdot\left(a'_{u,1}\right)^{\mathrm{wt}(i,j)} \tag{8.2}$$

for the following choice of weights $\mathrm{wt}(i,j)$, $(i,j)\neq(u,1)$. We define

$$\mathrm{wt}(i,j) = \Lambda_{\mathcal{M},\mathcal{W},\gamma,k}(i,j) \tag{8.3}$$

for

$$(i,j)\in\{(i,1)\mid i\in[1,k]\}\cup\{(i,2)\mid i\in[1,\gamma-1]\setminus\mathcal{M}\}.$$

In particular, by Definition 6.6 this gives $\mathrm{wt}(\gamma,1) = 1$ and $\mathrm{wt}(u,1) = 1$ although we don't mean to use $\mathrm{wt}(u,1)$ in (8.2). Also, we define $a'_{k,2} = a_{k,2} = 1$. By condition (x) the weight of any non-trivial Laurent monomial appearing on the right hand side of (8.1) equals -1.

Put

$$V' = \{a'_{i,1}\mid i\in[1,k]\}\cup\{a'_{i,2}\mid i\in[1,k-1]\setminus\mathcal{M}\}.$$

Define a collection of variables $W'_{\mathcal{M},\mathcal{W},\gamma,k}$ and a function

$$\Lambda'_{\mathcal{M},\mathcal{W},\gamma,k}\colon W'_{\mathcal{M},\mathcal{W},\gamma,k}\to\mathbb{Z}$$

replacing the variables $a_{i,j}$ by $a'_{i,j}$ in the appropriately applied Definition 6.6.

We rewrite (8.1) as

$$\frac{(1+a'_{k-1,1})\cdot\ldots\cdot(1+a'_{k-1,1}+\ldots+a'_{\gamma+1,1})}{a'_{k-1,1}\cdot\ldots\cdot a'_{\gamma+1,1}}\cdot\frac{1}{a'_{\gamma,1}\cdot a'_{u,1}} = \tag{8.4}$$

$$= 1 - \frac{1}{a'_{u,1}}\cdot\frac{P'}{M'} = \frac{M'\cdot a'_{u,1} - P'}{M'\cdot a'_{u,1}}.$$

Here P' is a polynomial that does not depend on the variables $a'_{u,1}$ and $a'_{k,1}$, and M' is a monomial that does not depend on the variables $a'_{u,1}$ and $a'_{k,1}$. Note that equation (8.4) is indeed valid even in the case when $\gamma = k$ (which means $u\neq k$) since in this case one has

$$\frac{(1+a'_{k-1,1})\cdot\ldots\cdot(1+a'_{k-1,1}+\ldots+a'_{\gamma+1,1})}{a'_{k-1,1}\cdot\ldots\cdot a'_{\gamma+1,1}} = 1.$$

We rewrite (8.4) as

$$a'_{\gamma,1} = \frac{M'}{M' \cdot a'_{u,1} - P'} \cdot \frac{(1 + a'_{k-1,1}) \cdot \ldots \cdot (1 + a'_{k-1,1} + \ldots + a'_{\gamma+1,1})}{a'_{k-1,1} \cdot \ldots \cdot a'_{\gamma+1,1}}. \tag{8.5}$$

Now we put

$$a''_{u,1} = \frac{M' \cdot a'_{u,1} - P'}{M'} \cdot \frac{a'_{k-1,1} \cdot \ldots \cdot a'_{\gamma+1,1}}{(1 + a'_{k-1,1}) \cdot \ldots \cdot (1 + a'_{k-1,1} + \ldots + a'_{\gamma+1,1})}, \tag{8.6}$$

we put $a''_{i,1} = a'_{i,1}$ for all $i \in [1, k-1]$, $i \neq u$, and we put $a''_{i,2} = a'_{i,2}$ for all $i \in [1, k] \setminus \mathcal{M}$ (in particular, this gives $a''_{k,2} = a'_{k,2} = 1$). Then

$$a_{u,1} = a'_{u,1} = a''_{u,1} \cdot \frac{(1 + a''_{k-1,1}) \cdot \ldots \cdot (1 + a''_{k-1,1} + \ldots + a''_{\gamma+1,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{\gamma+1,1}} + \frac{P''}{M''}, \tag{8.7}$$

where P'' and M'' are obtained from P' and M' by replacing the variables $a'_{i,j}$ by the corresponding variables $a''_{i,j}$. We can rewrite (8.5) as

$$\begin{aligned} a_{\gamma,1} = a'_{\gamma,1} \cdot a'_{u,1} = \\ = \frac{1}{a''_{u,1}} \cdot \left(a''_{u,1} \cdot \frac{(1 + a''_{k-1,1}) \cdot \ldots \cdot (1 + a''_{k-1,1} + \ldots + a''_{\gamma+1,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{\gamma+1,1}} + \frac{P''}{M''} \right). \end{aligned} \tag{8.8}$$

Also, (8.2) implies that

$$\begin{aligned} a_{i,j} = a'_{i,j} \cdot \left(a'_{u,1}\right)^{\mathrm{wt}(i,j)} = \\ = a''_{i,j} \cdot \left(a''_{u,1} \cdot \frac{(1 + a''_{k-1,1}) \cdot \ldots \cdot (1 + a''_{k-1,1} + \ldots + a''_{\gamma+1,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{\gamma+1,1}} + \frac{P''}{M''} \right)^{\mathrm{wt}(i,j)} \end{aligned} \tag{8.9}$$

for

$$(i,j) \in \{(i,1) \mid i \in [1,k], i \neq u, i \neq \gamma\} \cup \{(i,2) \mid i \in [1, \gamma-1] \setminus \mathcal{M}\}.$$

Equation (8.8) allows us to exclude the variable $a_{\gamma,1}$. Put

$$V'' = \{a''_{i,1} \mid i \in [1,k], i \neq \gamma\} \cup \{a''_{i,2} \mid i \in [1, \gamma-1] \setminus \mathcal{M}\}.$$

We define

$$\psi \colon \mathbb{T}(V'') \dashrightarrow \mathbb{T}(V)$$

to be the change of variables from $a_{i,j}$ to $a''_{i,j}$. We define the quiver $\mathcal{Q}''$ so that $\mathrm{Ver}(\mathcal{Q}'') = \mathrm{Ver}(\mathcal{Q})$, and the set $\mathrm{Ar}(\mathcal{Q}'')$ consists of a single arrow

$$\alpha_f = \langle (k,2) \to (k,3) \rangle.$$

Finally, we put $R''(i,j) = \psi^* R(i,j)$.

One has

$$\psi^* F_{\mathcal{Q},V,R} = \psi^* \left(\sum_{\alpha \in \mathrm{Ar}(\mathcal{Q})} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} \right) =$$

$$= \psi^* \left(\sum_{\alpha \in B} \frac{R\left(h(\alpha)\right)}{R\left(t(\alpha)\right)} + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} \right) =$$

$$= 1 + \frac{R\left(h(\alpha_f)\right)}{R\left(t(\alpha_f)\right)} = 1 + \psi^* R(k,3).$$

By condition (x) the rational function $R(k,3)$ is a Laurent polynomial in the variables of V such that the weight of any Laurent monomial of $R(k,3)$ is non-negative. Therefore, plugging (8.7), (8.8) and (8.9) into $R(k,3)$, we see that $\psi^* R(k,3)$, and thus also $\psi^* F_{\mathcal{Q},V,R}$, is a Laurent polynomials in the variables of V''. □

Remark 8.2 (cf. Remark 6.4). If in Lemma 8.1 we replace the equation $1 - F_{\mathcal{Q},V,R,B} = 0$ by $1 - F_{\mathcal{Q},V,R,B} = U$, then equation 8.5 takes the form

$$a'_{\gamma,1} = \frac{M'}{(1-U) \cdot M' \cdot a'_{u,1} - P'} \cdot \frac{(1 + a'_{k-1,1}) \cdot \ldots \cdot (1 + a'_{k-1,1} + \ldots + a'_{\gamma+1,1})}{a'_{k-1,1} \cdot \ldots \cdot a'_{\gamma+1,1}}. \tag{8.10}$$

We conclude this section by a couple of general remarks concerning the proofs of Lemmas 6.3, 6.8, 6.10, 7.1, 7.3 and 8.1.

Remark 8.3. Let $n = 2$, let $(\mathcal{Q}, V, R)$ be a triplet, and let $B \subset \mathrm{Ar}(\mathcal{Q})$ be a block such that a quiver with the set of vertices coinciding with $\mathrm{Ver}(\mathcal{Q})$ and the set of arrows $\mathrm{Ar}(\mathcal{Q}) \setminus B$ is admissible.

Suppose that $(\mathcal{Q}, V, R)$ is a result of an application of Lemma 6.3 to a triplet $(\widehat{\mathcal{Q}}, \widehat{V}, \widehat{R})$ and some horizontal block $\widehat{B} \subset \mathrm{Ar}(\widehat{\mathcal{Q}})$ with the properties described in the assumptions of Lemma 6.3. Then $(\mathcal{Q}, V, R)$ and B satisfy the assumptions of Lemmas 6.8, 6.10, 7.3 or 8.1 depending on whether B is a horizontal block of size at least 2, a basic horizontal block, a mixed block or a vertical block.

Similarly, suppose that $(\mathcal{Q}, V, R)$ is a result of an application of Lemma 6.8 to some triplet $(\widehat{\mathcal{Q}}, \widehat{V}, \widehat{R})$ and some horizontal block $\widehat{B} \subset \mathrm{Ar}(\widehat{\mathcal{Q}})$. Then $(\mathcal{Q}, V, R)$ and B satisfy the assumptions of Lemmas 6.8, 6.10, 7.3 or 8.1 depending on whether B is a horizontal block of size at least 2, a basic horizontal block, a mixed block or a vertical block.

Finally, suppose that $(\mathcal{Q}, V, R)$ is a result of an application of Lemma 6.10. Then $(\mathcal{Q}, V, R)$ and B satisfy the assumptions of Lemmas 6.10, 7.3 or 8.1 depending on whether B is a basic horizontal block, a mixed block or a vertical block. Note that we are not able to apply Lemma 6.8 to a result of an application of Lemma 6.10.

We will use these observations during the inductive proof of Theorem 5.3 below.

Remark 8.4. A reader may have an impression that our choice of main variables and weight variables in the proofs of Lemmas 6.3, 6.8, 7.1, 7.3 and 8.1 is rather arbitrary. This is true to some extent, and some choices could be maid in some other way. Nevertheless, at least part of our choices is inevitable, and some of the others are done due to our attempts to optimize the computations. First, when we choose a main variable for some block we want that the corresponding vertex is not simultaneously a tail of some arrow of the block and a head of some other arrow of the block. Thus one of the very few unnecessary

things here is the choice of the variable a instead of $a_{1,2}$ as a main variable in the proof of Lemma 6.3. We did this because we wanted to unify the case when the size of the block equals 1 and the case when the size of the block exceeds 1, and also, more importantly, to obtain a bit more uniform set of variables after this first change.

Furthermore, the weight of a weight variable with respect to itself is 1, and our method of expressing a main variable requires that for any arrow α of the block one has $\mathrm{wt}(t(\alpha)) = \mathrm{wt}(h(\alpha))+1$. Therefore, when working with a horizontal block B in the proof of Lemma 6.8 we choose a weight variable in the second line from below in $\mathrm{Ver}(B)$; this allows us to leave the variables corresponding to the last row of $\mathrm{Ver}(B)$ unaffected by the change of coordinates, so that the further changes of coordinates remain relatively simple, and so that we do not have a contradiction with assigning the weight to $a_{k,2} = 1$ if $(k, 2) \in \mathrm{Ver}(B)$. Also, in this case we choose a weight variable in the first column rather than in the second column of $\mathrm{Ver}(B)$ to avoid dealing with more cases that would arise if the block B could contain an arrow between the vertices corresponding to a main variable and a weight variable.

Similarly to this, when we choose a weight variable in the proof of Lemma 7.1, we choose it in the second column to avoid dealing with more cases that would arise if there could be an arrow in the block B between the vertices corresponding to the main variable a and a weight variable. Besides this, we are forced to choose the weight variable in the $(k-1)$-th row since the weight of $a_{k,2} = 1$ with respect to anything is 0.

In the proof of Lemma 7.3 our main variable corresponds, as explained above, to the unique vertex $(i, j) \in \mathrm{Ver}(B)$ that is not simultaneously a tail of some arrow of the block and a head of some other arrow of B, and such that the corresponding rational function $R(i, j)$ is a variable. On the other hand, the choice of the weight variable is dictated by the requirement that the distance between the corresponding vertex and the vertex $(k, 2) \in \mathrm{Ver}(B)$ along the arrows of B should equal 1. This leaves us with a choice between the variables $a_{k,1}$ and $a_{k-1,2}$, with no big difference between these cases.

Finally, in the proof of Lemma 8.1 we choose $a_{\gamma,1}$ to be the main variable since it is the only one that we actually managed to express via the remaining variables in the most general case. The choice of the weight variable $a_{u,1}$ is mostly defined by the function $\Lambda_{\mathcal{M},\mathcal{W},\gamma,k}$. In principle, u can be replaced by any number from the set $\mathcal{W}$, or by any number from the set $[\gamma + 1, k]$.

Remark 8.5. Our proofs of Lemmas 7.3 and 8.1 rely on different degree conditions (cf. condition (ix) of Lemma 7.3 and condition (x) of Lemma 8.1). We did not manage to unify them, but we suspect that it may be possible if one uses some other degree function.

9. Proof of the main theorem

In this section we prove Theorem 5.3 using preliminary computations performed in Sections 6, 7 and 8.

Proof of Theorem 5.3. Define an auxiliary triplet $(\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ as follows. Put $\widetilde{\mathcal{Q}}_0 = \mathcal{Q}_0$ and

$$\widetilde{V}_0 = \{\widetilde{a}_{i,1} \mid i \in [1, k]\} \cup \{\widetilde{a}_{i,2} \mid i \in [1, k-1]\} \cup \{a\}.$$

Define

$$\widetilde{R}_0(k, 2) = 1, \quad \widetilde{R}_0(0, 1) = \widetilde{R}_0(k, 3) = a,$$

and $\widetilde{R}_0(i,j) = \widetilde{a}_{i,j}$ for $i \in [1,k]$, $j \in [1,2]$, $(i,j) \neq (k,2)$. Let

$$\widetilde{\psi}_0 \colon \mathbb{T}(\widetilde{V}_0) \to \mathbb{T}(V_0)$$

be a monomial change of variables given by

$$\widetilde{a}_{i,j} = \frac{a_{i,j}}{a_{k,n}}, \quad a = \frac{1}{a_{k,n}}.$$

It is easy to check that

$$\widetilde{\psi}_0^*(F_{\mathcal{Q}_0,V_0,R_0}) = F_{\widetilde{\mathcal{Q}}_0,\widetilde{V}_0,\widetilde{R}_0}.$$

We choose the blocks $B_1, \ldots, B_l$ in the following way.

If $\sum d_i \leqslant k$, then we consecutively choose $B_1, \ldots, B_l$ to be horizontal blocks of size $d_1, \ldots, d_l$ situated as high as possible.

If $\sum d_i = k+1$ and $d_l \geqslant 2$, then we consecutively choose $B_1, \ldots, B_{l-1}$ to be horizontal blocks of size $d_1, \ldots, d_{l-1}$ situated as high as possible. After this we choose B_l to be a mixed block of size d_l, so that B_l covers all the remaining vertical arrows of $\mathrm{Ar}(\mathcal{Q}_0)$, and all horizontal arrows of $\mathrm{Ar}(\mathcal{Q}_0)$ except for the arrow $\langle (k,2) \to (k,3) \rangle$. In particular, if $l = 1$ and $d_1 = k+1$, then we choose B_1 to be the mixed block that consists of all arrows of $\mathrm{Ar}(\mathcal{Q}_0)$ except for the arrow $\langle (k,2) \to (k,3) \rangle$.

Finally, if $\sum d_i = k+1$ and $d_l = 1$, then we choose $B_1, \ldots, B_{l-1}$ to be horizontal blocks of size $d_1, \ldots, d_{l-1}$ situated as high as possible in the quiver $\mathcal{Q}_0$. This means that the union $B_1 \cup \ldots \cup B_{l-1}$ covers all vertical arrows of $\mathrm{Ar}(\mathcal{Q}_0)$. After this we choose B_l to be the first basic vertical block.

In other words, we always choose $B_1, \ldots, B_l$ so that the blocks B_i and B_j are disjoint for any $i \neq j$, $i,j \in [1,l]$, and for any $i \in [1,l]$ the quiver $\mathcal{Q}_i$ with $\mathrm{Ver}(\mathcal{Q}_i) = \mathrm{Ver}(\mathcal{Q}_0)$ and

$$\mathrm{Ar}(\mathcal{Q}_i) = \mathrm{Ar}(\mathcal{Q}_0) \setminus (B_1 \cup \ldots \cup B_i)$$

is admissible. Note also that the arrow $\langle (k,2) \to (k,3) \rangle$ is not contained in any of the blocks B_i.

We proceed to define the rational maps ψ_i. If B_1 is a mixed block define $\widetilde{\psi}_1$ by Lemma 7.1 and put $\psi_1 = \widetilde{\psi}_1 \circ \widetilde{\psi}_0$; in this case ψ_1 is the only change of variables we need.

If B_1 is a horizontal block define $\widetilde{\psi}_1$ by Lemma 6.3. Put $\psi_1 = \widetilde{\psi}_1 \circ \widetilde{\psi}_0$. We put $\mathcal{W}_1 = \varnothing$ and $\gamma_1 = 1$ if $d_1 = 1$, and we put $\mathcal{W}_1 = \{w\}$ and $\gamma_1 = w+1$, where $a_{w,1}$ is the weight variable used in the proof of Lemma 6.3, if $d_1 > 1$; we also put $\mathcal{M}_1 = \varnothing$. Note that $(\mathcal{M}_1, \mathcal{W}_1, \gamma_1)$ is a block history of $(\mathcal{Q}_1, \gamma_1)$. Then we consider the remaining horizontal blocks B_i one by one and define changes of variables ψ_i and block histories $(\mathcal{M}_i, \mathcal{W}_i, \gamma_i)$ by Lemmas 6.8 or 6.10, depending on whether the size of B_i exceeds 1 or equals 1. Note that due to our choice of the blocks we will first have to apply Lemma 6.8 several times, and then Lemma 6.10 several times. If $\sum d_i \leqslant k$, then this is all we need. If $\sum d_i = k+1$, then we conclude with a construction of ψ_l applying Lemma 7.3 or Lemma 8.1, depending on whether the block B_l is mixed or vertical.

Our final observation is that in the process described above we can always perform the next required step due to compatibility of conditions and assertions of Lemmas 6.3, 6.8, 7.3 and 8.1 pointed out in Remark 8.3. $\square$

10. Periods

In this section we check that Givental's integral gives the so called main period for complete intersections in projective spaces and Grassmannians of planes. To do this we start from an integral of the form of the left hand side of (3.9) over an indefinite cycle δ_1 (that we will specify later). Then we take residues several times obtaining integrals over cycles δ_i such that δ_{i-1} is a boundary of a tubular neighborhood of δ_i. After taking all residues we define a cycle we integrate over and define all other cycles one by one. It turns out that the cycle δ_1 we recover in this way is homologous to a standard cycle δ_1^0 we used in (3.9).

Definition 10.1. Let f be a Laurent polynomial in m variables $x_1, \dots, x_m$. Let $\Omega(x_1, \dots, x_m)$ be a standard logarithmic form defined in (3.5). The integral

$$I_f(t) = \int\limits_{|x_i|=\varepsilon_i} \frac{\Omega(x_1, \dots, x_m)}{1 - tf} = \sum_{j=0}^{\infty} t^j \cdot \int\limits_{|x_i|=\varepsilon_i} f^j \Omega(x_1, \dots, x_m) \in \mathbb{C}[[t]]$$

is called *the main period* for f, where ε_i are arbitrary positive numbers.

Remark 10.2. Let ϕ_j be a constant term of f^j. Then $I_f(t) = \sum \phi_j t^j$.

The following theorem (which is a mathematical folklore, see [Prz08, Proposition 2.3] or [CCGGK12, Theorem 3.2] for the proof) justifies this definition.

Theorem 10.3. *Let f be a Laurent polynomial in m variables. Let P be a Picard–Fuchs differential operator for a pencil of hypersurfaces in a torus provided by f. Then one has $P[I_f(t)] = 0$.*

Consider a smooth complete intersection $Y \subset \mathbb{P}^N$ of hypersurfaces of degrees $d_1, \dots, d_l$. Denote

$$d_0 = N + 1 - \sum d_i.$$

Assume that $d_0 \geqslant 1$, that is Y is a Fano variety. Anticanonical Givental's Landau–Ginzburg model for Y is given in a torus

$$(\mathbb{C}^*)^N \cong \operatorname{Spec} \mathbb{C}[a_{i,j}^{\pm 1}, y_s^{\pm 1}], \quad i \in [1, l], j \in [1, d_i], s \in [1, d_0 - 1],$$

by equations

$$a_{i,1} + \ldots + a_{i,d_i} = 1, \quad i \in [1, l], \tag{10.1}$$

with superpotential

$$w = y_1 + \ldots + y_{d_0 - 1} + \frac{1}{\prod a_{i,j} \prod y_i}.$$

The subvariety cut out by equations (10.1) after change of variables given by

$$x_{i,j} = \frac{a_{i,j}}{a_{i,d_i}}, \quad i \in [1, l], j \in [1, d_i - 1]$$

is birational to a torus

$$(\mathbb{C}^*)^m \cong \operatorname{Spec} \mathbb{C}[x_{i,j}^{\pm 1}, y_s^{\pm 1}], \quad i \in [1, l], j \in [1, d_i - 1], s \in [1, d_0 - 1],$$

where $m = N - l$. The superpotential w gives a Laurent polynomial

$$f_Y = \frac{\prod_{i=1}^{l} (x_{i,1} + \ldots + x_{i,d_i - 1} + 1)^{d_i}}{\prod_{i=1}^{l} \prod_{j=1}^{d_i - 1} x_{i,j} \prod_{j=1}^{d_0 - 1} y_j} + y_1 + \ldots + y_{d_0 - 1}$$

which is a toric Landau–Ginzburg model for Y, see [Prz13, §3.2] and [ILP13].

Proposition 10.4. *One has*

$$I_Y = \int\limits_{\substack{|x_{i,j}|=\varepsilon_{i,j}\\ |y_s|=\varepsilon_s}} \frac{\Omega(x_{1,1},\dots,x_{l,d_l-1},y_1,\dots,y_{d_0-1})}{1-tf_Y}.$$

Proof. Consider an integral

$$I = \int_{\delta_1} \frac{\Omega(a_{1,1},\dots,a_{l,d_l},y_1,\dots,y_{d_0-1})}{\prod_{i=1}^{l}(1-(a_{i,1}+\ldots+a_{i,d_i}))\cdot\left(1-t\cdot\left(\frac{1}{\prod a_{i,j}\prod y_s}+\sum y_s\right)\right)}$$

for some N-cycle δ_1, cf. (3.9).

Put

$$x_{i,j} = \frac{a_{i,j}}{a_{i,d_i}}, \quad i\in[1,l], j\in[1,d_i-1].$$

Then one has

$$I = \int_{\delta_1'} \frac{\pm\Omega(x_{1,1},\dots,x_{1,d_1-1},\dots,x_{l,1},\dots,x_{l,d_l-1},a_{1,d_1},\dots,a_{l,d_l},y_1,\dots,y_{d_0-1})}{\prod_{i=1}^{l}\left(1-\left(\sum_{j=1}^{d_i-1}x_{i,j}+1\right)\cdot a_{i,d_i}\right)\cdot\left(1-t\cdot\left(\frac{1}{\prod x_{i,j}\prod a_{i,d_i}^{d_i}\prod y_s}+\sum y_s\right)\right)}$$

for some N-cycle δ_1'.

Finally put

$$Q_i = 1-\left(\sum_{j=1}^{d_i-1}x_{i,j}+1\right)\cdot a_{i,d_i}, \quad i\in[1,l],$$

so that

$$a_{i,d_i} = \frac{1-Q_i}{\sum_{j=1}^{d_i-1}x_{i,j}+1}.$$

After this we have

$$I = \int_{\delta_1''} \frac{\pm\Omega(x_{1,1},\dots,x_{l,d_l-1},Q_1,\dots,Q_l,y_1,\dots,y_{d_0-1})}{\prod_{i=1}^{l}(1-Q_i)\cdot\left(1-t\cdot\left(\frac{\prod_{i=1}^{l}(x_{i,1}+\ldots+x_{i,d_i-1}+1)^{d_i}}{\prod_{i=1}^{l}(1-Q_i)^l\prod_{j=1}^{d_i-1}x_{i,j}\prod_{s=1}^{d_0-1}y_s}+y_1+\ldots+y_{d_0-1}\right)\right)}$$

for some N-cycle δ_1''. Taking residues with respect to variables Q_i, possibly reordering and renaming variables one gets

$$I = \int_{\Delta} \frac{\Omega(x_{1,1},\dots,x_{l,d_l-1},y_1,\dots,y_{d_0-1})}{1-tf_Y}$$

for some m-cycle Δ.

Put $\Delta = \{|x_{i,j}| = \varepsilon_{i,j}, |y_s| = \varepsilon_s\}$ and define cycles $\delta_2,\dots,\delta_{l+1} = \Delta$ so that δ_{i-1} is a boundary of a tubular neighborhood of δ_i for $i\in[3,l+1]$, and δ_1'' is a boundary of a tubular neighborhood of δ_2. One can check that δ_1'', and thus also δ_1 is homologous to a cycle

$$\delta_1^0 = \{|a_{i,j}| = \varepsilon_{i,j}, |y_s| = \varepsilon_s\}$$

which completes the proof. □

Now we check that Givental's integral gives the main period for complete intersections in Grassmannians of planes as well.

Proposition 10.5. *Consider a smooth Fano complete intersection*

$$Y = \mathrm{Gr}(2, k+2) \cap Y_1 \cap \ldots \cap Y_l$$

and a nef-partition corresponding to the choice of blocks from the proof of Theorem 5.3. Let I_Y^0 be Givental's integral for this nef-partition, and $\widehat{f}_Y = F_{\mathcal{Q}_l, V_l, R_l}$ be a Laurent polynomial in $m = 2k - l$ variables $x_1, \ldots, x_m$ given by Theorem 5.3. Put $f_Y = \widehat{f}_Y - l$. Then

$$I_Y^0 = \int\limits_{|x_i|=\varepsilon_i} \frac{\Omega(x_1, \ldots, x_m)}{1 - t f_Y}.$$

Proof. We choose blocks $B_1, \ldots, B_l$ as in the proof of Theorem 5.3 and put

$$B_0 = \mathrm{Ar}(\mathcal{Q}_0) \setminus \left(\cup_{i \in [1,l]} B_i\right).$$

Note that one has $f_Y = \psi^* F_{\mathcal{Q}_0, V_0, R_0, B_0}$ where $\psi \colon \mathbb{T}(V_l) \dashrightarrow \mathbb{T}(V_0)$ is the change of variables given by Theorem 5.3. Consider an integral

$$I = \int_{\delta_1} \frac{\Omega(a_{i,j})}{\prod_{s=1}^{l} (1 - F_{\mathcal{Q}_0, V_0, R_0, B_i}) \cdot (1 - t F_{\mathcal{Q}_0, V_0, R_0, B_0})}$$

over an indefinite $2k$-cycle δ_1, cf. (3.9). Applying the monomial change variables described in the beginning of the proof of Theorem 5.3 we obtain an integral

$$I = \int_{\delta_1'} \frac{\pm \Omega(a, \widetilde{a}_{i,j})}{\prod_{s=1}^{l} (1 - F_{\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0, B_i}) \cdot (1 - t F_{\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0, B_0})}$$

for some $2k$-cycle δ_1'.

We follow changes of variables from the proof of Theorem 5.3. Consider a form

$$\Omega = \frac{\Omega(a_1, \ldots, a_p) \cdot F(a_1, \ldots, a_p)}{1 - F_{\mathcal{Q}, V, R, B}},$$

where $F_{\mathcal{Q}, V, R, B}$ depends on some variables a_i and a function $F(a_1, \ldots, a_p)$ is chosen so that

$$I = \int_{\delta_j} \frac{\Omega}{1 - t f}$$

for some Laurent polynomial f and some p-cycle δ_j. Denote $1 - F_{\mathcal{Q}, V, R, B}$ by U. Depending on whether the block B is a horizontal block containing the arrow $\langle (0,1) \to (1,1) \rangle$, a horizontal block of size at least 2 not containing the arrow $\langle (0,1) \to (1,1) \rangle$, a basic horizontal block not containing the arrow $\langle (0,1) \to (1,1) \rangle$, a mixed block containing the arrow $\langle (0,1) \to (1,1) \rangle$, a mixed block not containing the arrow $\langle (0,1) \to (1,1) \rangle$, or the first basic vertical block, we follow changes of variables described in Lemmas 6.3, 6.8, 6.10, 7.1, 7.3, or 8.1 respectively.

Suppose that we are not in the situation described in Lemma 6.10. Let a_1 be a main variable and a_2 be a weight variable for the change of variables for the change of variables that agrees with $(\mathcal{Q}, V, R)$ and B. The latter can be decomposed in several changes of variables. The first change of variables is monomial so by equation (3.7) a standard logarithmic form in new variables is equal to a standard logarithmic form in the initial variables a_i. If we are in the situation described in Lemma 6.10, then we do not change variables on this step and choose a_2 to be the variable $a_{r,1}$ in the notation of Lemma 6.10.

Keeping the same notation for changed variables for simplicity the form Ω can be written down as

$$\frac{da_1}{a_1}\wedge\frac{da_2}{a_2}\wedge\frac{\Omega(a_3,\dots,a_p)\cdot F(a_1,\dots,a_p)}{U}.$$

We put

$$a_1=\frac{T}{(1-U)\cdot a_2-S}$$

for certain Laurent polynomials S and T in $a_3,\dots,a_p$, like in (6.16), (6.33), (6.38), (7.11), (7.25), and (8.10).

After the this substitution the form Ω is

$$\frac{da_1}{a_1}\wedge\frac{da_2}{a_2}\wedge\frac{\Omega(a_3,\dots,a_p)\cdot F(a_1,\dots,a_p)}{U}=$$

$$=\big((1-U)\cdot a_2-S\big)\cdot d\left(\frac{1}{(1-U)\cdot a_2-S}\right)\wedge\frac{da_2}{a_2}\wedge\frac{\Omega(a_3,\dots,a_p)\cdot F\left(\frac{1}{(1-U)a_2-S},a_2,\dots,a_p\right)}{U}=$$

$$=\frac{-1}{(1-U)\cdot a_2-S}\cdot\frac{dU}{U}\wedge da_2\wedge\Omega(a_3,\dots,a_p)\cdot F\left(\frac{1}{(1-U)a_2-S},a_2,\dots,a_p\right).$$

After taking a residue with respect to U we get

$$\operatorname{Res}_U\Omega=\frac{-1}{a_2-S}\cdot da_2\wedge\Omega(a_3,\dots,a_p)\cdot F\left(\frac{1}{a_2-S},a_2,\dots,a_p\right).$$

We put

$$a_2=R\cdot b+S$$

for some Laurent polynomial R in $a_3,\dots,a_p$, like in (6.9), (6.24), (6.35), (7.8), (7.19), and (8.7). Now our new variables are $b,a_3,\dots,a_p$. After this substitution we get

$$\operatorname{Res}_U\Omega=\frac{-1}{a_2-S}\cdot da_2\wedge\Omega(a_3,\dots,a_p)\cdot F\left(\frac{1}{a_2-S},a_2,\dots,a_p\right)=$$

$$=\frac{-1}{R\cdot b}\cdot d\left(R\cdot b+S\right)\wedge\Omega(a_3,\dots,a_p)\cdot F\left(\frac{1}{R\cdot b},R\cdot b+S,a_3\dots,a_p\right)=$$

$$=-\frac{db}{b}\wedge\Omega(a_3,\dots,a_p)\cdot F\left(\frac{1}{R\cdot b},R\cdot b+S,a_3\dots,a_p\right).$$

Thus

$$I=\int_{\delta_j}\frac{\Omega}{1-tf}=\int_{\delta_{j+1}}-\Omega(b,a_3,\dots,a_p)\cdot\overline{F}\left(b,a_3\dots,a_p\right)$$

for some $(p-1)$-cycle δ_{j+1}, where

$$\overline{F}\left(b,a_3\dots,a_p\right)=F\left(\frac{1}{R\cdot b},R\cdot b+S,a_3\dots,a_p\right).$$

Applying this procedure step by step l times following the proof of Theorem 5.3, we define cycles $\delta_2,\dots,\delta_{l+1}=\Delta$ so that δ_{i-1} is a boundary of a tubular neighborhood of δ_i for $i\in[3,l+1]$, and δ_1' is a boundary of a tubular neighborhood of δ_2 and arrive to an integral

$$\int\limits_{\Delta}\frac{\Omega(x_1,\dots,x_m)}{1-tf_Y}$$

for some Laurent polynomial f_Y in some variables $x_1, \dots, x_m$. Put $\Delta = \{|x_i| = \varepsilon_i\}$ and recover the cycles $\delta_1', \delta_2, \dots, \delta_l$. One can check that δ_1', and thus also δ_1 is homologous to a cycle

$$\delta_1^0 = \{|a_{i,j}| = \varepsilon_{i,j}\}$$

which completes the proof. □

Corollary 10.6. *The proof of Theorem 5.3 provides very weak Landau–Ginzburg models for complete intersections in Grassmannians of planes.*

Proof. Let Y be a complete intersection in a Grassmannian of planes and let f_Y be a Laurent polynomial given by Theorem 5.3. In other words a family of hypersurfaces in a torus corresponding to f_Y is relatively birational to anticanonical Givental's Landau–Ginzburg model for Y. By [BCFKS98] and [BCFK03, Proposition 3.5] Givental's integral for (Y, ω_Y), where ω_Y is an anticanonical form, equals $\widetilde{I}^Y$. On the other hand, by Remark 10.2 it is a constant terms series of f_Y, i. e. a main period of Y. □

11. Hyperplane sections

In this section we apply Theorem 5.3 to obtain explicit formulas for Laurent polynomials corresponding to Fano varieties that are sections of Grassmannians of planes by several hyperplanes. We will use notation introduced in Theorem 5.3. Keeping in mind Remark 3.4 and Proposition 10.5, we will be more interested in the shifted Laurent polynomials $F_{\mathcal{Q}_l, V_l, R_l} - l$ than $F_{\mathcal{Q}_l, V_l, R_l}$ themselves.

Lemma 11.1. *Suppose that $n = 2$, $l \leqslant k$ and $d_1 = \ldots = d_l = 1$. Consider the triplet $(\mathcal{Q}_0, V_0, R_0)$. Let B_i, $i \in [1, l]$, be the $(i-1)$-th basic horizontal block. Then there is a sequence of triplets $(\mathcal{Q}_i, V_i, R_i)$, $i \in [1, l]$, and a sequence of changes of variables*

$$\psi_i \colon \mathbb{T}(V_i) \dashrightarrow \mathbb{T}(V_{i-1}), \quad i \in [1, l],$$

such that the change of variables ψ_i agrees with the triplet $(\mathcal{Q}_{i-1}, V_{i-1}, R_{i-1})$ and the block B_i, the triplet $(\mathcal{Q}_i, V_i, R_i)$ is a transformation of the triplet $(\mathcal{Q}_{i-1}, V_{i-1}, R_{i-1})$ associated to ψ_i, one has

$$V_l = \{a_{i,1} \mid i \in [1, k]\} \cup \{a_{i,2} \mid i \in [l, k-1]\},$$

and the following conditions hold:

(i) *the quiver $\mathcal{Q}_l$ does not contain vertical arrows α such that $h(\alpha) = (i, j)$ for $i \in [1, l]$, $j \in [1, 2]$;*
(ii) *one has $R(k, 2) = 1$;*
(iii) *for (i, j) with $i \in [l, k]$, $j \in [1, 2]$, $(i, j) \neq (k, 2)$, one has $R_l(i, j) = a_{i,j}$;*
(iv) *for any $i \in [1, l-1]$ one has*

$$R_l(i, 1) = a_{l,1} + a_{l-1,1} + \ldots + a_{i,1};$$

(v) *for any $i \in [1, l-1]$ one has*

$$R_l(i, 2) = \frac{a_{l,2} \cdot (a_{l,1} + a_{l-1,1}) \cdot (a_{l,1} + a_{l-1,1} + a_{l-2,1}) \cdot \ldots \cdot (a_{l,1} + a_{l-1,1} + \ldots + a_{i,1})}{a_{l-1,1} \cdot a_{l-2,1} \cdot \ldots \cdot a_{i,1}}$$

if $l < k$, and

$$R_k(i, 2) = \frac{(a_{k,1} + a_{k-1,1}) \cdot (a_{k,1} + a_{k-1,1} + a_{k-2,1}) \cdot \ldots \cdot (a_{k,1} + a_{k-1,1} + \ldots + a_{i,1})}{a_{k-1,1} \cdot a_{k-2,1} \cdot \ldots \cdot a_{i,1}}$$

if $l = k$.

(vi) *one has*

$$R_l(k,3) = R(1,1) = a_{l,1} + a_{l-1,1} + \ldots + a_{1,1}.$$

Proof. Arguing as in the proof of Theorem 5.3, we start with the standard triplet $(\mathcal{Q}_0, V_0, R_0)$ and obtain a triplet $(\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ as described in the proof of Theorem 5.3. Then we apply Lemma 6.3 with $s = 1$ once, and apply Lemma 6.10 with $\gamma = 1$ and $\mathcal{M} = \mathcal{W} = \varnothing$ consecutively $l - 1$ times. □

Applying Lemma 11.1, we immediately obtain.

Corollary 11.2. *In the notation of Lemma 11.1 suppose that* $l \leqslant k - 1$. *Then one has*

$$\begin{aligned} (11.1)\quad & F_{\mathcal{Q}_l, V_l, R_l} - l = \\ &= \sum_{i \in [1, l-1]} \frac{a_{l,2} \cdot (a_{l,1} + a_{l-1,1}) \cdot \ldots \cdot (a_{l,1} + \ldots + a_{i+1,1})}{a_{l-1,1} \cdot \ldots \cdot a_{i,1}} + \sum_{i \in [l, k-1]} \frac{a_{i,2}}{a_{i,1}} + \frac{1}{a_{k,1}} + \\ &\qquad + \sum_{\substack{i \in [l, k-2] \\ j \in [1,2]}} \frac{a_{i+1,j}}{a_{i,j}} + \frac{a_{k,1}}{a_{k-1,1}} + \frac{1}{a_{k-1,2}} + a_{l,1} + a_{l-1,1} + \ldots + a_{1,1}. \end{aligned}$$

Corollary 11.3. *In the notation of Lemma 11.1 suppose that* $l = k$. *Then one has*

$$\begin{aligned} (11.2)\quad & F_{\mathcal{Q}_k, V_k, R_k} - k = \\ &= \sum_{i \in [1, k-1]} \frac{(a_{k,1} + a_{k-1,1}) \cdot \ldots \cdot (a_{k,1} + \ldots + a_{i+1,1})}{a_{k-1,1} \cdot \ldots \cdot a_{i,1}} + \frac{1}{a_{k,1}} + a_{k,1} + a_{k-1,1} + \ldots + a_{1,1}. \end{aligned}$$

Now we proceed to the case corresponding to a Fano variety that is a section of the Grassmannian $\mathrm{Gr}(2, k+2)$ by $k+1$ hyperplanes.

Lemma 11.4. *Suppose that* $n = 2$, $l = k + 1$ *and* $d_1 = \ldots = d_{k+1} = 1$. *Consider the triplet* $(\mathcal{Q}_0, V_0, R_0)$. *Let* B_i, $i \in [1, k]$, *be the* $(i-1)$*-th basic horizontal block, and let* B_{k+1} *be the first basic vertical block. Then there is a sequence of triplets* $(\mathcal{Q}_i, V_i, R_i)$, $i \in [1, k+1]$, *and a sequence of changes of variables*

$$\psi_i \colon \mathbb{T}(V_i) \dashrightarrow \mathbb{T}(V_{i-1}), \quad i \in [1, k+1],$$

such that the change of variables ψ_i *agrees with the triplet* $(\mathcal{Q}_{i-1}, V_{i-1}, R_{i-1})$ *and the block* B_i, *the triplet* $(\mathcal{Q}_i, V_i, R_i)$ *is a transformation of the triplet* $(\mathcal{Q}_{i-1}, V_{i-1}, R_{i-1})$ *associated to* ψ_i, *one has*

$$V_{k+1} = \{a_{i,1} \mid i \in [2, k]\}$$

and

$$F_{\mathcal{Q}_{k+1},V_{k+1},R_{k+1}} - (k+1) = \tag{11.3}$$
$$= \left(a_{k,1} \cdot \frac{(1+a_{k-1,1}) \cdot \ldots \cdot (1+a_{k-1,1}+\ldots+a_{2,1})}{a_{k-1,1} \cdot \ldots \cdot a_{2,1}} + \right.$$
$$\left. + \sum_{i \in [2,k-1]} \frac{(1+a_{k-1,1}) \cdot \ldots \cdot (1+\ldots+a_{i+1,1})}{a_{k-1,1} \cdot \ldots \cdot a_{i,1}} + 1 \right) \times$$
$$\times \left(1 + a_{k-1,1} + \ldots + a_{2,1} + \frac{1}{a_{k,1}} \right).$$

Proof. We obtain changes of variables $\psi_1, \ldots, \psi_k$ from Lemma 11.1, and proceed to make a change of variables ψ_{k+1} that agrees with the triplet $(\mathcal{Q}_k, V_k, R_k)$ and the block B_{k-1}.

To do this we apply Lemma 8.1 with $\gamma = 1$ and $\mathcal{M} = \mathcal{W} = \varnothing$. As prescribed in the proof of Lemma 8.1, we choose $a_{k,1}$ to be the weight variable and $a_{1,1}$ to be the main variable. We put $\mathrm{wt}(i,1) = 1$ for $i \in [1,k]$, rewrite (8.7) as

$$a_{k,1} = a''_{k,1} \cdot \frac{(1+a''_{k-1,1}) \cdot \ldots \cdot (1+a''_{k-1,1}+\ldots+a''_{2,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{2,1}} + \tag{11.4}$$
$$+ \sum_{i \in [2,k-1]} \frac{(1+a''_{k-1,1}) \cdot \ldots \cdot (1+\ldots+a''_{i+1,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{i,1}} + 1,$$

and rewrite (8.8) as

$$a_{1,1} = \frac{1}{a''_{k,1}} \cdot \left(a''_{k,1} \cdot \frac{(1+a''_{k-1,1}) \cdot \ldots \cdot (1+a''_{k-1,1}+\ldots+a''_{2,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{2,1}} + \right. \tag{11.5}$$
$$\left. + \sum_{i \in [1,k-1]} \frac{(1+a''_{k-1,1}) \cdot \ldots \cdot (1+\ldots+a''_{i+1,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{i,1}} + 1 \right).$$

Also, by (8.9) we have

$$a_{i,1} = a''_{i,1} \cdot \left(a''_{k,1} \cdot \frac{(1+a''_{k-1,1}) \cdot \ldots \cdot (1+a''_{k-1,1}+\ldots+a''_{2,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{2,1}} + \right. \tag{11.6}$$
$$\left. + \sum_{i \in [1,k-1]} \frac{(1+a''_{k-1,1}) \cdot \ldots \cdot (1+\ldots+a''_{i+1,1})}{a''_{k-1,1} \cdot \ldots \cdot a''_{i,1}} + 1 \right)$$

for any $i \in [2, k-1]$.

We arrive to a triplet $(\mathcal{Q}_{k+1}, V_{k+1}, R_{k+1})$ such that $\mathrm{Ar}(\mathcal{Q}_{k+1})$ consists of a single arrow $\langle (k,2) \to (k,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_{k+1} = \{ a_{i,1} \mid i \in [2,k] \}.$$

Now we obtain equation (11.3) by a direct computation using (11.4), (11.5) and (11.6). $\square$

Remark 11.5 (cf. Problem 13.3). One can easily see that families of hypersurfaces given by equations (11.1), (11.2) and (11.3) can be compactified to singular Calabi–Yau hypersurfaces by multiplying by denominators and homogenizing.

Problem 11.6 (cf. Problem 13.4)**.** *Prove that the compactifications mentioned in Remark 11.5 admit crepant resolutions. In other words, prove that these very weak Landau–Ginzburg models are weak ones. In addition prove that the corresponding toric varieties admit smoothings to hyperplane sections of Grassmannians, that is, prove that equations* (11.1)*,* (11.2) *and* (11.3) *give toric Landau–Ginzburg models.*

12. Examples

In this section we provide several sporadic examples that illustrate our computations performed in Sections 6, 7 and 8. We will use notation introduced in Theorem 5.3. Keeping in mind Remark 3.4 and Proposition 10.5, we will be more interested in the shifted Laurent polynomials $F_{\mathcal{Q}_l,V_l,R_l} - l$ than $F_{\mathcal{Q}_l,V_l,R_l}$ themselves, just as in Section 11. Very weak Landau–Ginzburg models for threefold examples (coinciding with ours up to monomial changes of variables) can be found in [Prz13] and [CCGGK12], where they are obtained by different methods.



Figure 6. Triplets $(\mathcal{Q}_0, V_0, R_0)$ and $(\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ for the Grassmannian $\mathrm{Gr}(2,4)$

Example 12.1. The following computation corresponds to a quadric threefold, which we treat as a hyperplane section of the Grassmannian $\mathrm{Gr}(2,4)$.

Let $k = n = 2$, $l = 1$ and $d_1 = 1$ in the notation of Theorem 5.3. In this case equation (11.1) gives

$$F_{\mathcal{Q}_1,V_1,R_1} - 1 = \frac{a_{1,2}}{a_{1,1}} + \frac{1}{a_{2,1}} + \frac{a_{2,1}}{a_{1,1}} + \frac{1}{a_{1,2}} + a_{1,1}.$$

This polynomial is, up to monomial change of variables, the toric Landau–Ginzburg model for quadric threefold written down in [KP12, Example 2.2].

Example 12.2. The following computation corresponds to a Fano fourfold of index 3 that is a section of the Grassmannian $\mathrm{Gr}(2,5)$ by two hyperplanes.

Let $n = 2$, $k = 3$, $l = 2$ and $d_1 = d_2 = 1$. In this case equation (11.1) gives

$$F_{\mathcal{Q}_2,V_2,R_2} - 2 = \frac{a_{2,2}}{a_{1,1}} + \frac{a_{2,2}}{a_{2,1}} + \frac{1}{a_{3,1}} + \frac{a_{3,1}}{a_{2,1}} + \frac{1}{a_{2,2}} + a_{2,1} + a_{1,1}.$$

Example 12.3. The following computation corresponds to a smooth quadric surface, which we treat as an intersection of two hyperplanes in the Grassmannian $\mathrm{Gr}(2,4)$. The same result was known earlier; actually, it is just a simplified Givental's Landau–Ginzburg model for the quadric surface treated as a toric variety $\mathbb{P}^1 \times \mathbb{P}^1$, see Section 3.

Let $n = k = 2$, $l = 2$ and $d_1 = d_2 = 1$. In this case (11.2) gives

$$F_{\mathcal{Q}_2,V_2,R_2} - 2 = \frac{1}{a_{1,1}} + \frac{1}{a_{2,1}} + a_{2,1} + a_{1,1}.$$

Example 12.4. The following example provides another computation that corresponds to a smooth quadric surface. Similarly to Example 12.3, we treat the quadric surface as an intersection of two hyperplanes in the Grassmannian $\mathrm{Gr}(2,4)$, and we use variable changes that agree with various blocks to obtain the result, but unlike Example 12.3 (or rather Lemma 11.1 where the actual computation is performed) we do not follow exactly the procedure prescribed by the proof of Theorem 5.3. Our point here is that our procedure is not the only one, and sometimes not even the shortest one, to obtain the answer.

We have $n = k = 2$. We start with the standard triplet $(\mathcal{Q}_0, V_0, R_0)$ and obtain a triplet $(\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ as described in the proof of Theorem 5.3, see Figure 6. Then we make variable changes that agree with the 0-th horizontal basic block, which consists of a single arrow $\langle (0,1) \to (1,1) \rangle$, and with the second vertical basic block, which consists of a single arrow $\langle (2,2) \to (2,3) \rangle$. This gives us two equations $\widetilde{a} = \widetilde{a}_{1,1}$ and $\widetilde{a} = 1$, which we use to exclude variables $\widetilde{a}$ and $\widetilde{a}_{1,1}$. This gives us a triplet $(\mathcal{Q}, V, R)$ such that $\mathrm{Ar}(\mathcal{Q})$ consists of the arrows $\langle (1,1) \to (1,2) \rangle$, $\langle (2,1) \to (2,2) \rangle$, $\langle (1,1) \to (2,1) \rangle$ and $\langle (1,2) \to (2,2) \rangle$, one has $V = \{a_{1,2}, a_{2,1}\}$, and

$$R(0,1) = R(1,1) = R(2,2) = R(2,3) = 1, \quad R(1,2) = a_{1,2}, \quad R(2,1) = a_{2,1}.$$

We compute

$$F_{\mathcal{Q},V,R} - 2 = a_{1,2} + a_{2,1} + \frac{1}{a_{1,2}} + \frac{1}{a_{2,1}}.$$

More than this, one can start from the triplet $(\mathcal{Q}_0, V_0, R_0)$ itself, and utilize the same two blocks to obtain equations $a_{1,1} = 1$ and $a_{2,2} = 1$. Using these equations to exclude variables $a_{1,1}$ and $a_{2,2}$ we obtain exactly the same result as above.

Example 12.5. The following computation corresponds to a Fano threefold of anti-canonical degree 40 and index 2 that is a section of the Grassmannian $\mathrm{Gr}(2,5)$ by three hyperplanes (see e. g. [IP99, §3.4]). Due to [Ga07], this variety has a terminal Gorenstein toric degeneration. One can see that the Laurent polynomial that we get is given by a procedure discussed in Section 3. In fact it is a toric Landau–Ginzburg model, see [Prz13, Theorem 18].

Let $n = 2$, $k = 3$, $l = 3$ and $d_1 = d_2 = d_3 = 1$. In this case (11.2) gives

$$F_{\mathcal{Q}_3,V_3,R_3} - 3 = \frac{a_{3,1} + a_{2,1}}{a_{2,1} \cdot a_{1,1}} + \frac{1}{a_{2,1}} + \frac{1}{a_{3,1}} + a_{3,1} + a_{2,1} + a_{1,1}.$$

Example 12.6. The following computation corresponds to a Fano fourfold of index 2 that is a section of the Grassmannian $\mathrm{Gr}(2,6)$ by four hyperplanes.

Let $n = 2$, $k = 4$, $l = 4$ and $d_1 = d_2 = d_3 = d_4 = 1$. In this case equation (11.2) gives

$$F_{\mathcal{Q}_4,V_4,R_4} - 4 = \\ = \frac{(a_{4,1} + a_{3,1}) \cdot (a_{4,1} + a_{3,1} + a_{2,1})}{a_{3,1} \cdot a_{2,1} \cdot a_{1,1}} + \frac{a_{4,1} + a_{3,1}}{a_{3,1} \cdot a_{2,1}} + \frac{1}{a_{3,1}} + \frac{1}{a_{4,1}} + a_{4,1} + a_{3,1} + a_{2,1} + a_{1,1}.$$

In [PSh14] the relative compactification of a family of hypersurfaces in $(\mathbb{C}^*)^4$ given by this Laurent polynomial is computed. This computation confirms expectations of Homological Mirror Symmetry in this case.

Example 12.7. The following computation corresponds to a Fano fivefold of index 2 that is a section of the Grassmannian $\mathrm{Gr}(2, 7)$ by five hyperplanes.

Let $n = 2$, $k = 5$, $l = 5$ and $d_1 = d_2 = d_3 = d_4 = d_5 = 1$. In this case (11.2) gives

$$F_{\mathcal{Q}_5,V_5,R_5} - 5 = \frac{(a_{5,1} + a_{4,1}) \cdot (a_{5,1} + a_{4,1} + a_{3,1}) \cdot (a_{5,1} + a_{4,1} + a_{3,1} + a_{2,1})}{a_{4,1} \cdot a_{3,1} \cdot a_{2,1} \cdot a_{1,1}} + \\ + \frac{(a_{5,1} + a_{4,1}) \cdot (a_{5,1} + a_{4,1} + a_{3,1})}{a_{4,1} \cdot a_{3,1} \cdot a_{2,1}} + \frac{a_{5,1} + a_{4,1}}{a_{4,1} \cdot a_{3,1}} + \frac{1}{a_{4,1}} + \frac{1}{a_{5,1}} + \\ + a_{5,1} + a_{4,1} + a_{3,1} + a_{2,1} + a_{1,1}.$$

Example 12.8. The following computation corresponds to a del Pezzo surface of degree 5 that is a section of the Grassmannian $\mathrm{Gr}(2, 5)$ by four hyperplanes.

Let $n = 2$, $k = 3$, $l = 4$ and $d_1 = d_2 = d_3 = d_4 = 1$. In this case equation (11.3) gives

$$F_{\mathcal{Q}_4,V_4,R_4} - 4 = \left(a_{3,1} \cdot \frac{1 + a_{2,1}}{a_{2,1}} + \frac{1}{a_{2,1}} + 1 \right) \cdot \left(1 + a_{2,1} + \frac{1}{a_{3,1}} \right).$$

Example 12.9. The following computation corresponds to a Fano threefold of anti-canonical degree 14 and index 1 that is a section of the Grassmannian $\mathrm{Gr}(2, 6)$ by five hyperplanes (see [IP99, §12.2]).

Let $n = 2$, $k = 4$, $l = 5$ and $d_1 = d_2 = d_3 = d_4 = d_5 = 1$. In this case equation (11.3) gives

$$F_{\mathcal{Q}_5,V_5,R_5} - 5 = \\ = \left(a_{4,1} \cdot \frac{(1 + a_{3,1}) \cdot (1 + a_{3,1} + a_{2,1})}{a_{3,1} \cdot a_{2,1}} + \frac{1 + a_{3,1}}{a_{3,1} \cdot a_{2,1}} + \frac{1}{a_{3,1}} + 1 \right) \times \\ \times \left(1 + a_{2,1} + a_{3,1} + \frac{1}{a_{4,1}} \right).$$

Example 12.10. The following computation corresponds to a three-dimensional complete intersection of two quadrics (one of which we treat as the Grassmannian $\mathrm{Gr}(2, 4)$).

Let $n = k = 2$, $l = 1$ and $d_1 = 2$. We start with the standard triplet $(\mathcal{Q}_0, V_0, R_0)$ and obtain a triplet $(\mathcal{Q}, V, R) = (\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ as described in the proof of Theorem 5.3, see Figure 6. Simplifying the notation we have

$$V = \{a, a_{1,1}, a_{1,2}, a_{2,1}\},$$

and

$$R(0, 1) = R(2, 3) = a, \quad R(1, 1) = a_{1,1}, \quad R(1, 2) = a_{1,2}, \quad R(2, 1) = a_{2,1}, \quad R(2, 2) = 1.$$

Now we proceed to make a change of variables ψ that agrees with the triplet $(\mathcal{Q}, V, R)$ and the mixed block B that consists of all vertical arrows of $\mathrm{Ar}(\mathcal{Q})$, i. e. of the arrows $\langle (0,1) \to (1,1) \rangle$, $\langle (1,1) \to (2,1) \rangle$ and $\langle (1,2) \to (2,2) \rangle$. As prescribed in the proof of Lemma 6.3, we choose $a_{1,1}$ to be the weight variable, and we choose a to be the main variable. We find that $\delta' = 1$ in the notation of (6.6), rewrite (6.9) as

$$a_{1,1} = a''_{1,1} + \frac{a''_{1,2} \cdot a''_{2,1} + 1}{a''_{1,2}},$$

and rewrite (6.13) as

$$a = \frac{1}{a''_{1,1}} \cdot \left(a''_{1,1} + \frac{a''_{1,2} \cdot a''_{2,1} + 1}{a''_{1,2}} \right)^2.$$

We arrive to a triplet $(\mathcal{Q}_1, V_1, R_1)$ such that $\mathrm{Ar}(\mathcal{Q}_1)$ consists of all horizontal arrows of $\mathrm{Ar}(\mathcal{Q})$, i. e. of the arrows $\langle (1,1) \to (1,2) \rangle$, $\langle (2,1) \to (2,2) \rangle$ and $\langle (2,2) \to (2,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_1 = \{a_{1,1}, a_{1,2}, a_{2,1}\}.$$

We also have

$$R_1(0,1) = R_1(2,3) = \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + a_{2,1} + \frac{1}{a_{1,2}} \right)^2, \quad R_1(1,1) = a_{1,1} + a_{2,1} + \frac{1}{a_{1,2}}, \tag{12.1}$$

$$R_1(1,2) = a_{1,2} \cdot \left(a_{1,1} + a_{2,1} + \frac{1}{a_{1,2}} \right), \quad R_1(2,1) = a_{2,1}, \quad R_1(2,2) = 1.$$

Finally, we compute

$$F_{\mathcal{Q}_1, V_1, R_1} - 1 = a_{1,2} + \frac{1}{a_{2,1}} + \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + a_{2,1} + \frac{1}{a_{1,2}} \right)^2.$$

Example 12.11. The following computation corresponds to a two-dimensional complete intersection of two quadrics, which we treat as a section of the Grassmannian $\mathrm{Gr}(2,4)$ by a quadric and a hypersurface.

Let $n = k = 2$, $l = 2$, $d_1 = 2$ and $d_2 = 1$. Arguing as in Example 12.10, we make a variable change that agrees with the horizontal block B_1 of size 2 and arrive to a triple $(\mathcal{Q}_1, V_1, R_1)$ such that $\mathrm{Ar}(\mathcal{Q}_1)$ consists of all horizontal arrows of $\mathrm{Ar}(\mathcal{Q}_0)$, one has

$$V_1 = \{a_{1,1}, a_{1,2}, a_{2,1}\},$$

and the rational functions $R_1(i,j)$ are described by (12.1).

We proceed to make a change of variables ψ_2 that agrees with the triplet $(\mathcal{Q}_1, V_1, R_1)$ and the first basic vertical block B_2. To do this we apply Lemma 8.1 with $\gamma = 2$, $\mathcal{M} = \varnothing$ and $\mathcal{W} = \{1\}$. As prescribed in the proof of Lemma 8.1, we choose $a_{1,1}$ to be the weight variable and $a_{2,1}$ to be the main variable. We put

$$\mathrm{wt}(1,1) = \mathrm{wt}(2,1) = 1, \quad \mathrm{wt}(1,2) = -1,$$

rewrite (8.7) as

$$a_{1,1} = a''_{1,1} + a''_{1,2},$$

and rewrite (8.8) as

$$a_{2,1} = \frac{a''_{1,1} + a''_{1,2}}{a''_{1,1}}.$$

Also, by (8.9) we have

$$a_{1,2} = \frac{a''_{1,2}}{a''_{1,1} + a''_{1,2}}.$$

We arrive to a triplet $(\mathcal{Q}_2, V_2, R_2)$ such that $\mathrm{Ar}(\mathcal{Q}_2)$ consists of a single arrow $\langle (2,2) \to (2,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_2 = \{a_{1,1}, a_{1,2}\}.$$

We also have $R_2(2,2) = 1$ and

$$F_{\mathcal{Q}_2, V_2, R_2} - 2 = R_2(2,3) = (a_{1,1} + a_{1,2}) \cdot \left(1 + \frac{1}{a_{1,1}} + \frac{1}{a_{1,2}}\right)^2.$$

Example 12.12. The following computation corresponds to a three-dimensional complete intersection of a quadric (which we treat as the Grassmannian $\mathrm{Gr}(2,4)$) and a cubic.

Let $n = k = 2$, $l = 1$ and $d_1 = 3$. We start with the standard triplet $(\mathcal{Q}_0, V_0, R_0)$ and obtain a triplet $(\mathcal{Q}, V, R) = (\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ as described in the proof of Theorem 5.3, see Figure 6. Simplifying the notation we have

$$V = \{a, a_{1,1}, a_{1,2}, a_{2,1}\},$$

and

$$R(0,1) = R(2,3) = a, \quad R(1,1) = a_{1,1}, \quad R(1,2) = a_{1,2}, \quad R(2,1) = a_{2,1}, \quad R(2,2) = 1,$$

exactly as in Example 12.10.

Now we proceed to make a change of variables ψ that agrees with the triplet $(\mathcal{Q}, V, R)$ and the mixed block B that consists of all arrows of $\mathrm{Ar}(\mathcal{Q})$ except for the arrow $\langle (2,2) \to (2,3) \rangle$. As prescribed in the proof of Lemma 7.1, we choose $a_{1,2}$ to be the weight variable, and we choose a to be the main variable. We find that $\Delta' = 1$ in the notation of (7.4), rewrite (7.8) as

$$a_{1,2} = a''_{1,2} + \frac{(a''_{2,1})^2 + a''_{1,1} \cdot a''_{2,1} + a''_{1,1} + a''_{2,1}}{a''_{1,1} \cdot a''_{2,1}},$$

and rewrite (7.10) as

$$a = \frac{a''_{1,1}}{a''_{1,2}} \cdot \left(a''_{1,2} + \frac{(a''_{2,1})^2 + a''_{1,1} \cdot a''_{2,1} + a''_{1,1} + a''_{2,1}}{a''_{1,1} \cdot a''_{2,1}}\right)^3.$$

We arrive to a triplet $(\mathcal{Q}_1, V_1, R_1)$ such that $\mathrm{Ar}(\mathcal{Q}_1)$ consists of a single arrow $\langle (2,2) \to (2,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_1 = \{a_{1,1}, a_{1,2}, a_{2,1}\}.$$

We also have $R_1(2,2) = 1$ and

$$F_{\mathcal{Q}_1, V_1, R_1} - 1 = R_1(2,3) = \frac{a_{1,1}}{a_{1,2}} \cdot \left(a_{1,2} + \frac{a_{2,1}^2 + a_{1,1} \cdot a_{2,1} + a_{1,1} + a_{2,1}}{a_{1,1} \cdot a_{2,1}}\right)^3.$$

Example 12.13. The following computation corresponds to a Fano fourfold of anticanonical degree 160 and index 2 that is a section of the Grassmannian $\mathrm{Gr}(2,5)$ by a quadric and a hyperplane (see e. g. [DIM12]).

Let $n = 2$, $k = 3$, $l = 2$, $d_1 = 2$ and $d_2 = 1$. We start with the standard triplet $(\mathcal{Q}_0, V_0, R_0)$ and obtain a triplet $(\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ as described in the proof of Theorem 5.3, see Figures 4 and 5. Simplifying the notation for variables, at this step we have

$$\widetilde{V}_0 = \{a, a_{1,1}, a_{1,2}, a_{2,1}, a_{2,2}, a_{3,1}\},$$

and

$$\widetilde{R}_0(0,1) = R_1(3,3) = a, \quad \widetilde{R}_0(3,2) = 1,$$
$$\widetilde{R}_0(i,j) = a_{i,j} \text{ for } (i,j) \in \{(1,1),(1,2),(2,1),(2,2),(3,1)\}.$$

We proceed to make a change of variables $\tilde{\psi}_1$ that agrees with the triplet $(\widetilde{\mathcal{Q}}_0, \widetilde{V}_0, \widetilde{R}_0)$ and the horizontal block B_1. As prescribed in the proof of Lemma 6.3, we choose $a_{1,1}$ to be the weight variable and a to be the main variable, rewrite (6.9) as

$$a_{1,1} = a''_{1,1} + a''_{2,1} + \frac{a''_{2,2}}{a''_{1,2}},$$

and rewrite (6.13) as

$$a = \frac{1}{a''_{1,1}} \cdot \left(a''_{1,1} + a''_{2,1} + \frac{a''_{2,2}}{a''_{1,2}} \right)^2.$$

We arrive to a triplet $(\mathcal{Q}_1, V_1, R_1)$ such that $\mathrm{Ar}(\mathcal{Q}_1)$ consists of all arrows of $\mathrm{Ar}(\mathcal{Q}_0)$ except for $\langle (0,1) \to (1,1) \rangle$, $\langle (1,1) \to (2,1) \rangle$ and $\langle (1,2) \to (2,2) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_1 = \{a_{1,1}, a_{1,2}, a_{2,1}, a_{2,2}, a_{3,1}\},$$

and

$$R_1(0,1) = R_1(3,3) = \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + a_{2,1} + \frac{a_{2,2}}{a_{1,2}} \right)^2, \tag{12.2}$$
$$R_1(1,1) = a_{1,1} + a_{2,1} + \frac{a_{2,2}}{a_{1,2}}, \quad R_1(1,2) = a_{1,2} \cdot \left(a_{1,1} + a_{2,1} + \frac{a_{2,2}}{a_{1,2}} \right),$$
$$R_1(3,2) = 1, \quad R_1(i,j) = a_{i,j} \text{ for } (i,j) \in \{(2,1),(2,2),(3,1)\}.$$

We also have

$$F_{\mathcal{Q}_1, V_1, R_1} - 1 = \frac{a_{3,1}}{a_{2,1}} + \frac{1}{a_{2,2}} + a_{1,2} + \frac{a_{2,2}}{a_{2,1}} + \frac{1}{a_{3,1}} + \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + a_{2,1} + \frac{a_{2,2}}{a_{1,2}} \right)^2.$$

We proceed to make a change of variables ψ_2 that agrees with the triplet $(\mathcal{Q}_1, V_1, R_1)$ and the second horizontal basic block B_2. To do this we apply Lemma 6.10 with $\gamma = 2$, $\mathcal{M} = \varnothing$ and $\mathcal{W} = \{1\}$. As prescribed in the proof of Lemma 6.10, we choose $a_{2,2}$ to be the main variable, rewrite (6.35) as

$$a_{2,1} = a'_{2,1} + a'_{3,1},$$

and rewrite (6.36) as

$$a_{2,2} = \frac{a'_{2,1} + a'_{3,1}}{a'_{2,1}}.$$

We arrive to a triplet $(\mathcal{Q}_2, V_2, R_2)$ such that $\mathrm{Ar}(\mathcal{Q}_2)$ consists of all horizontal arrows of $\mathrm{Ar}(\mathcal{Q}_0)$, i. e. of the arrows $\langle (1,1) \to (1,2) \rangle$, $\langle (2,1) \to (2,2) \rangle$, $\langle (3,1) \to (3,2) \rangle$

and $\langle (3,2) \to (3,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_2 = \{a_{1,1}, a_{1,2}, a_{2,1}, a_{3,1}\},$$

and

$$R_2(0,1) = R_2(3,3) = \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + a_{2,1} + a_{3,1} + \frac{1}{a_{1,2}} + \frac{a_{3,1}}{a_{1,2} \cdot a_{2,1}} \right)^2, \tag{12.3}$$

$$R_2(1,1) = a_{1,1} + a_{2,1} + a_{3,1} + \frac{1}{a_{1,2}} + \frac{a_{3,1}}{a_{1,2} \cdot a_{2,1}},$$

$$R_2(1,2) = a_{1,2} \cdot \left(a_{1,1} + a_{2,1} + a_{3,1} + \frac{1}{a_{1,2}} + \frac{a_{3,1}}{a_{1,2} \cdot a_{2,1}} \right),$$

$$R_2(2,1) = a_{2,1} + a_{3,1}, \quad R_2(2,2) = \frac{a_{2,1} + a_{3,1}}{a_{2,1}}, \quad R_2(3,1) = a_{3,1}, \quad R_2(3,2) = 1.$$

Finally, we have

$$F_{\mathcal{Q}_2, V_2, R_2} - 2 = a_{1,2} + \frac{1}{a_{2,1}} + \frac{1}{a_{3,1}} + \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + a_{2,1} + a_{3,1} + \frac{1}{a_{1,2}} + \frac{a_{3,1}}{a_{1,2} \cdot a_{2,1}} \right)^2.$$

Example 12.14. The following computation corresponds to a Fano threefold of anti-canonical degree 10 and index 1 that is an intersection of the Grassmannian $\mathrm{Gr}(2,5)$ with a quadric and two hyperplanes (see e. g. [IP99, §5.1], [DIM12]).

Let $n = 2$, $k = 3$, $l = 3$, $d_1 = 2$ and $d_2 = d_3 = 1$. Arguing as in Example 12.13, we make variable changes that agree with the horizontal blocks B_1 and B_2 and arrive to a triplet $(\mathcal{Q}_2, V_2, R_2)$ such that $\mathrm{Ar}(\mathcal{Q}_2)$ consists of all horizontal arrows of $\mathrm{Ar}(\mathcal{Q}_0)$, one has

$$V_2 = \{a_{1,1}, a_{1,2}, a_{2,1}, a_{3,1}\},$$

and the rational functions $R_2(i,j)$ are described by (12.3).

We proceed to make a change of variables ψ_3 that agrees with the triplet $(\mathcal{Q}_2, V_2, R_2)$ and the first vertical basic block B_3. To do this we apply Lemma 8.1 with $\gamma = 2$, $\mathcal{M} = \varnothing$ and $\mathcal{W} = \{1\}$. As prescribed in the proof of Lemma 8.1, we choose $a_{3,1}$ to be the weight variable and $a_{2,1}$ to be the main variable. We put

$$\mathrm{wt}(1,1) = \mathrm{wt}(2,1) = \mathrm{wt}(3,1) = 1, \quad \mathrm{wt}(1,2) = -1,$$

rewrite (8.7) as

$$a_{3,1} = a''_{3,1} + a''_{1,2} + 1,$$

and rewrite (8.8) as

$$a_{2,1} = \frac{a''_{3,1} + a''_{1,2} + 1}{a''_{3,1}}.$$

Also, by (8.9) we have

$$a_{1,1} = a''_{1,1} \cdot (a''_{3,1} + a''_{1,2} + 1)$$

and

$$a_{1,2} = \frac{a''_{1,2}}{a''_{3,1} + a''_{1,2} + 1}.$$

We arrive to a triplet $(\mathcal{Q}_3, V_3, R_3)$ such that $\mathrm{Ar}(\mathcal{Q}_3)$ consists of a single arrow $\langle (3,2) \to (3,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_3 = \{a_{1,1}, a_{1,2}, a_{3,1}\}.$$

Finally, we have

$$F_{\mathcal{Q}_3, V_3, R_3} - 3 = R_3(3,3) = \frac{a_{3,1} + a_{1,2} + 1}{a_{1,1}} \cdot \left(a_{1,1} + \frac{1}{a_{3,1}} + 1 + \frac{1}{a_{1,2}} + \frac{a_{3,1}}{a_{1,2}} \right)^2.$$

Example 12.15. The following computation corresponds to a Fano fourfold of anticanonical degree 20 and index 1 that is an intersection of the Grassmannian $\mathrm{Gr}(2,5)$ with two quadrics.

Let $n = 2$, $k = 3$, $l = 2$ and $d_1 = d_2 = 2$. Arguing as in the beginning of Example 12.13, we make a variable change that agrees with the horizontal block B_1 and arrive to a triplet $(\mathcal{Q}_1, V_1, R_1)$ such that $\mathrm{Ar}(\mathcal{Q}_1)$ consists of all arrows of $\mathrm{Ar}(\mathcal{Q}_0)$ except for the arrows $\langle (0,1) \to (1,1) \rangle$, $\langle (1,1) \to (2,1) \rangle$ and $\langle (1,2) \to (2,2) \rangle$, one has

$$V_1 = \{a_{1,1}, a_{1,2}, a_{2,1}, a_{2,2}, a_{3,1}\},$$

and the rational functions $R_1(i,j)$ are described by (12.2).

We proceed to make a change of variables ψ_2 that agrees with the triplet $(\mathcal{Q}_1, V_1, R_1)$ and the mixed block B_2 of size 2. To do this we apply Lemma 7.3 with $\mathcal{M} = \varnothing$ and $\mathcal{W} = \{1\}$. As prescribed in the proof of Lemma 7.3, we choose $a_{3,1}$ to be the weight variable and $a_{2,1}$ to be the main variable. We find that $\Delta' = \delta' = 1$ in the notation of (7.15), rewrite (7.19) as

$$a_{3,1} = a''_{3,1} + \frac{1 + a''_{1,2} \cdot a''_{2,2} + a''_{2,2}}{a''_{2,2}},$$

and rewrite (7.21) as

$$a_{2,1} = \frac{1 + a''_{2,2}}{a''_{3,1}} \cdot \left(a''_{3,1} + \frac{1 + a''_{1,2} \cdot a''_{2,2} + a''_{2,2}}{a''_{2,2}} \right)^2.$$

Also, due to the choice of the weights in the proof of Lemma 7.3 we have

$$a_{2,2} = a''_{2,2} \cdot \left(a''_{3,1} + \frac{1 + a''_{1,2} \cdot a''_{2,2} + a''_{2,2}}{a''_{2,2}} \right)$$

and

$$a_{1,2} = a''_{1,2} \cdot \left(a''_{3,1} + \frac{1 + a''_{1,2} \cdot a''_{2,2} + a''_{2,2}}{a''_{2,2}} \right)^{-1},$$

while $a_{1,1} = a''_{1,1}$.

We arrive to a triplet $(\mathcal{Q}_2, V_2, R_2)$ such that $\mathrm{Ar}(\mathcal{Q}_2)$ consists of a single arrow $\langle (3,2) \to (3,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_2 = \{a_{1,1}, a_{1,2}, a_{2,2}, a_{3,1}\}.$$

We also have $R_2(3,2)=1$ and

$$F_{\mathcal{Q}_2,V_2,R_2}-2=R_2(3,3)=$$
$$=\frac{1}{a_{1,1}}\cdot\left(a_{1,1}+\left(\frac{1+a_{2,2}}{a_{3,1}}+\frac{a_{2,2}}{a_{1,2}}\right)\cdot\left(a_{3,1}+\frac{1+a_{1,2}\cdot a_{2,2}+a_{2,2}}{a_{2,2}}\right)^2\right)^2.$$

Example 12.16. The following computation corresponds to a Fano fivefold of anticanonical degree 56 and index 1 that is an intersection of the Grassmannian $\mathrm{Gr}(2,6)$ with two quadrics and a hyperplane.

Let $n=2$, $k=4$, $l=3$ and $d_1=d_2=2$, $d_3=1$. We start with the standard triplet $(\mathcal{Q}_0,V_0,R_0)$, and after a change of variable corresponding to the horizontal block B_1 of size 2 we obtain a triplet $(\mathcal{Q}_1,V_1,R_1)$ as in the beginning of Example 12.13. Simplifying the notation for variables, at this step we have

$$V_1=\{a_{1,1},a_{1,2},a_{2,1},a_{2,2},a_{3,1},a_{3,2},a_{4,1}\},$$

and

$$R_1(0,1)=R_1(4,3)=\frac{1}{a_{1,1}}\cdot\left(a_{1,1}+a_{2,1}+\frac{a_{2,2}}{a_{1,2}}\right)^2,$$
$$R_1(1,1)=a_{1,1}+a_{2,1}+\frac{a_{2,2}}{a_{1,2}},\quad R_1(1,2)=a_{1,2}\cdot\left(a_{1,1}+a_{2,1}+\frac{a_{2,2}}{a_{1,2}}\right),$$
$$R_1(4,2)=1,\quad R_1(i,j)=a_{i,j}\text{ for }(i,j)\in\{(2,1),(2,2),(3,1),(3,2),(4,1)\}.$$

We also have

$$F_{\mathcal{Q}_1,V_1,R_1}-1=\frac{a_{3,1}}{a_{2,1}}+\frac{a_{3,2}}{a_{2,2}}+\frac{a_{4,1}}{a_{3,1}}+\frac{1}{a_{3,2}}+a_{1,2}+\frac{a_{2,2}}{a_{2,1}}+\frac{a_{3,2}}{a_{3,1}}+\frac{1}{a_{4,1}}+\frac{1}{a_{1,1}}\cdot\left(a_{1,1}+a_{2,1}+\frac{a_{2,2}}{a_{1,2}}\right)^2.$$

We proceed to make a change of variables ψ_2 that agrees with the triplet $(\mathcal{Q}_1,V_1,R_1)$ and the horizontal block B_2 of size 2. As prescribed in the proof of Lemma 6.8, we choose $a_{3,1}$ to be the weight variable and $a_{2,2}$ to be the main variable, rewrite (6.24) as

$$a_{3,1}=a''_{3,1}+\frac{a''_{3,2}+a''_{2,1}\cdot a''_{3,2}\cdot a''_{4,1}+a''_{2,1}}{a''_{2,1}\cdot a''_{3,2}},$$

and rewrite (6.27) as

$$a_{2,2}=\frac{a''_{3,2}}{a''_{3,1}}\cdot\left(a''_{3,1}+\frac{a''_{3,2}+a''_{2,1}\cdot a''_{3,2}\cdot a''_{4,1}+a''_{2,1}}{a''_{2,1}\cdot a''_{3,2}}\right)^2.$$

We arrive to a triplet $(\mathcal{Q}_2,V_2,R_2)$ such that $\mathrm{Ar}(\mathcal{Q}_2)$ consists of the arrows $\langle(1,1)\to(1,2)\rangle$, $\langle(2,1)\to(2,2)\rangle$, $\langle(3,1)\to(3,2)\rangle$, $\langle(4,1)\to(4,2)\rangle$ and $\langle(4,2)\to(4,3)\rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_2=\{a_{1,1},a_{1,2},a_{2,1},a_{3,1},a_{3,2},a_{4,1}\},$$

and

$$R_2(0,1) = R_2(4,3) =$$
$$= \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + \left(a_{2,1} + \frac{a_{3,2}}{a_{3,1} \cdot a_{1,2}} \right) \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right)^2 \right)^2,$$
$$R_2(1,1) = a_{1,1} + \left(a_{2,1} + \frac{a_{3,2}}{a_{3,1} \cdot a_{1,2}} \right) \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right)^2,$$
$$R_2(1,2) = a_{1,2} \cdot \left(a_{1,1} + \left(a_{2,1} + \frac{a_{3,2}}{a_{3,1} \cdot a_{1,2}} \right) \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right)^2 \right),$$
$$R_2(2,1) = a_{2,1} \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right)^2,$$
$$R_2(2,2) = \frac{a_{3,2}}{a_{3,1}} \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right)^2,$$
$$R_2(3,1) = a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}},$$
$$R_2(3,2) = a_{3,2} \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right),$$
$$R_2(4,1) = a_{4,1}, \quad R_2(4,2) = 1.$$

We also have

$$F_{\mathcal{Q}_2,V_2,R_2} - 2 = a_{1,2} + \frac{a_{3,2}}{a_{2,1} \cdot a_{3,1}} + a_{3,2} + \frac{1}{a_{4,1}} +$$
$$+ \frac{1}{a_{1,1}} \cdot \left(a_{1,1} + \left(a_{2,1} + \frac{a_{3,2}}{a_{3,1} \cdot a_{1,2}} \right) \cdot \left(a_{3,1} + \frac{a_{3,2} + a_{2,1} \cdot a_{3,2} \cdot a_{4,1} + a_{2,1}}{a_{2,1} \cdot a_{3,2}} \right)^2 \right)^2.$$

We proceed to make a change of variables ψ_3 that agrees with the triplet $(\mathcal{Q}_2, V_2, R_2)$ and the first vertical basic block B_3. To do this we apply Lemma 8.1 with $\gamma = 4$, $\mathcal{M} = \{2\}$ and $\mathcal{W} = \{1,3\}$. As prescribed in the proof of Lemma 8.1, we choose $a_{1,1}$ to be the weight variable and $a_{4,1}$ to be the main variable. We put

$$\mathrm{wt}(1,1) = \mathrm{wt}(3,1) = \mathrm{wt}(4,1) = 1, \quad \mathrm{wt}(1,2) = \mathrm{wt}(2,1) = \mathrm{wt}(3,2) = -1,$$

rewrite (8.7) as

$$a_{1,1} = a''_{1,1} + a''_{3,2} + \frac{a''_{3,2}}{a''_{2,1} \cdot a''_{3,1}} + a''_{1,2},$$

and rewrite (8.8) as

$$a_{4,1} = \frac{1}{a''_{1,1}} \cdot \left(a''_{1,1} + a''_{3,2} + \frac{a''_{3,2}}{a''_{2,1} \cdot a''_{3,1}} + a''_{1,2} \right).$$

Also, by (8.9) we have

$$a_{1,2} = a''_{1,2} \cdot \left(a''_{1,1} + a''_{3,2} + \frac{a''_{3,2}}{a''_{2,1} \cdot a''_{3,1}} + a''_{1,2} \right)^{-1},$$

$$a_{2,1} = a''_{2,1} \cdot \left(a''_{1,1} + a''_{3,2} + \frac{a''_{3,2}}{a''_{2,1} \cdot a''_{3,1}} + a''_{1,2} \right)^{-1},$$

$$a_{3,2} = a''_{3,2} \cdot \left(a''_{1,1} + a''_{3,2} + \frac{a''_{3,2}}{a''_{2,1} \cdot a''_{3,1}} + a''_{1,2} \right)^{-1}$$

and

$$a_{3,1} = a''_{3,1} \cdot \left(a''_{1,1} + a''_{3,2} + \frac{a''_{3,2}}{a''_{2,1} \cdot a''_{3,1}} + a''_{1,2} \right).$$

We arrive to a triplet $(\mathcal{Q}_3, V_3, R_3)$ such that $\mathrm{Ar}(\mathcal{Q}_3)$ consists of a single arrow $\langle (4,2) \to (4,3) \rangle$. Dropping for simplicity the primes from the notation for variables, we have

$$V_3 = \{a_{1,1}, a_{1,2}, a_{2,1}, a_{3,1}, a_{3,2}\}.$$

Finally, we compute

$$F_{\mathcal{Q}_3,V_3,R_3} - 3 = R_3(4,3) = \left(a_{1,1} + a_{1,2} + a_{3,2} + \frac{a_{3,2}}{a_{2,1} \cdot a_{3,1}} \right) \times$$
$$\times \left(1 + \left(a_{2,1} + \frac{a_{3,2}}{a_{3,1} \cdot a_{1,2}} \right) \cdot \left(a_{3,1} + \frac{1}{a_{2,1}} + \frac{1}{a_{3,2}} + \frac{1}{a_{1,1}} \right)^2 \right)^2.$$

Remark 12.17. Changes of variables described in Theorem 5.3 and choices of basic blocks for them are not unique ones that give Laurent polynomials. However in all cases we know all these Laurent polynomials for a given variety differ by cluster mutations, that is they correspond to families of hypersurfaces that are fiberwise birational and have a common (Calabi–Yau) compactification. It would be interesting to find out if this is true in general?

13. Discussion

Changes of variables discussed in sections 6, 7, 8, and 9 can be made for complete intersections in all Grassmannians $\mathrm{Gr}(n, k+n)$. Moreover, in [BCFKS98, Lemma 3.2.2] and [BCFKS98, Theorem 3.2.13] the natural generalization of maximal nef-partition from Grassmannians to partial flag varieties is suggested. So we expect that our proof of Theorem 5.3 can be generalized to these cases.

Problem 13.1 (see [DH15] and [PSh15b])**.** *Following Theorem 5.3 and Proposition 10.5 show the existence of very weak Landau–Ginzburg models for all complete intersections in Grassmannians and, more generally, partial flag varieties.*

Let us mention that to solve Problem 13.1 it is not enough to represent Givental's type Landau–Ginzburg models by Laurent polynomials. One should keep track of a particular type of change of variables in the spirit of Proposition 10.5 to check that under them the periods are preserved (cf. [PSh15b, Proposition 4.4]). However our experience shows that checking that a period condition is preserved for explicitly described birational transformations does not cause any difficulties.

In Theorem 5.3 we used a specific nef-partition to construct a Laurent polynomial. However sometimes one can use another nef-partitions and get a Laurent polynomial as well. In Example 12.4 we use other nef-partition to get the same result as one gets by Theorem 5.3. In some examples we consider one can get, using different nef-partitions, different Laurent polynomials. However they are mutationally equivalent to ones we get by Theorem 5.3.

Question 13.2. *Is this always the case?*

One more motivation for this question is given by [Li13]; the similar result is announced in [CKP14, Theorem 5.1] as a part of T. Prince's Thesis. That is, there are different methods that, under some assumptions, allow one to obtain Laurent polynomials for Givental's Landau–Ginzburg models. In [Li13] and in T. Prince's Thesis the resulting Laurent polynomials are proved to be actually independent on a choice of a nef-partition: Laurent polynomials obtained from different nef-partitions are relatively birational. In other words, they differ by mutations (cf. Example A.1).

According to a private communication with A. Harder, the families of hypersurfaces in tori given by Laurent polynomials we obtain in the proof of Theorem 5.3 have relative compactifications that are general complete intersections in toric varieties (cf. Remark 11.5). Moreover, they are Calabi–Yau compactifications. These compactifications enable one to compute the number of components of the unique reducible fibers of the compactifications.

Problem 13.3. *Prove the existence of Calabi–Yau compactifications of very weak Landau–Ginzburg models for complete intersections in Grassmannians of planes obtained in the proof of Theorem 5.3. In other words, prove that these models are weak ones. If Problem 13.1 is solved, prove this for complete intersections in arbitrary Grassmannians or, more generally, partial flag varieties.*

Another natural problem is the following.

Problem 13.4 (cf. [DH15])**.** *Prove that the toric varieties whose fan polytopes are Newton polytopes of very weak Landau–Ginzburg models for complete intersections in Grassmannians of planes obtained in the proof of Theorem 5.3 can be smoothed to corresponding complete intersections. In other words, prove that these models are toric ones. If Problem 13.1 is solved, prove this for complete intersections in arbitrary Grassmannians or, more generally, partial flag varieties.*

Problem 13.1 can be generalized further. A lot of interesting examples of higher-dimensional Fano varieties are not complete intersections in Grassmannians or partial flag manifolds but sections of non-decomposable vector bundles such as symmetric or skew powers of tautological vector bundle.

Question 13.5. *How to describe their analogs of nef-partitions for Grassmannians or partial flag varieties for smooth Fano varieties that are sections of non-decomposable vector bundles. Does the analog of Theorem 5.3 holds for them? Can analogs of Problems 13.3 and 13.4 be solved for them?*

Suppose that Problem 13.3 is solved. Let Y be a complete intersection of dimension r in Grassmannian of planes and let $LG(Y)$ be its Calabi–Yau compactification. Since birational smooth Calabi–Yau varieties are birational in codimension one, the number of

irreducible components in each fiber of $LG(Y)$ does not depend on a particular compactification. It is expected that there is at most one reducible fiber of $LG(Y)$. Denote the number of its irreducible components by $\widehat{k}$, and put $k_{LG(Y)} = \widehat{k} - 1$. This number can be computed via the approach mentioned above that was communicated to us by A. Harder.

Conjecture 13.6 (see [GKR12] and [PSh15a, Conjecture 1.1]). *Let $r \geqslant 3$. Then*

$$k_{LG(Y)} = h^{1,r-1}(Y).$$

For $r = 2$ one has

$$k_{LG(Y)} = h^{1,1}(Y) - 1.$$

Problem 13.7. *Prove Conjecture 13.6 for very weak Landau–Ginzburg models for complete intersections in Grassmannians of planes obtained in the proof of Theorem 5.3. If Problem 13.1 is solved, prove this for complete intersections in arbitrary Grassmannians or, more generally, partial flag varieties.*

Appendix A. Laurent polynomial mirrors for Grassmannians of planes, by Thomas Prince

In this appendix we describe an alternative approach to finding Laurent polynomial mirrors to complete intersections in Grassmannians. This calculation follows a general method for finding torus charts on Givental's Landau–Ginzburg model to complete intersections in toric varieties, detailed in [CKP14] generalizing the method of [Prz10]. In particular this method finds a torus chart on the Landau–Ginzburg model obtained by applying Givental's procedure to the flat toric degeneration of [BCFKS97]. In this appendix we shall restrict to Grassmannians of planes $\mathrm{Gr}(2, 2+k)$. We first recall the general form of this procedure.

Acknowledgements. The author is supported by an EPSRC Prize Studentship.

Following the observation made just after Theorem 4.1 that we may replace a complete intersection in $\mathrm{Gr}(2, 2+k)$ with one in $P(2, 2+k)$. We recall the general setup for Givental's procedure from Section 3, in the slightly more general setting which applies to complete intersections in $P(2, 2+k)$.

Assume we have fixed a Fano toric variety X, and a collection $L_1, \ldots, L_l$ of nef line bundles on X. As in Section 3 we let $\mathcal{N} \cong \mathbb{Z}^N$ denote the lattice containing the fan of X and $\mathcal{D} \cong \mathbb{Z}^{N+\rho}$ be a based lattice, with basis D_i, $i \in [1, N+\rho]$. Assuming the ray map is surjective we have the following exact sequence, see (3.2):

$$0 \longrightarrow \mathbb{L} \xrightarrow{D^*} \mathcal{D} \xrightarrow{r} \mathcal{N} \longrightarrow 0.$$

Defining $\mathbb{L}$ as the kernel, note that if X is smooth $\mathbb{L} = \operatorname{Pic} X^\vee$. We shall denote both the basis of $\mathcal{D}^\vee$ and the image of these basis elements in $\mathbb{L}^\vee$, under the dual map $D\colon \mathcal{D}^\vee \to \mathbb{L}^\vee$ by D_i, $i \in [1, N+\rho]$. We shall denote by D_i' the dual basis in $\mathcal{D}$. We shall assume that $(X; L_1, \ldots, L_l)$ satisfies the following conditions:

(1) X has a smooth zero-dimensional toric stratum. This determines a maximal-dimensional smooth cone in $\mathcal{N}$, and hence there is a subset $E \subset [1, N+\rho]$ such that the vectors $D_i, i \in E$ are a basis of $\mathbb{L}^\vee$. This condition implies that the ray map r is surjective.
(2) Each L_i is a non-negative combination of the $D_i, i \in E$.

(3) For each L_i, $i \in [1,l]$ there is a set $E_i \subset [1, N+\rho]$ such that $L_i = \sum_{j \in E_i} D_j$, and the collection of subsets E_k together with E is pairwise disjoint.

In other words we can define a *nef-partition* for $L_1, \ldots, L_l$ analogously to Section 3. We shall also fix an element $s_j \in E_j$ for each j and let $E_j^\circ = E_j \backslash \{s_j\}$. The map $D \colon \mathcal{D}^\vee \to \mathbb{L}^\vee$ is now a map of based lattices, and from now on we shall use D to refer to this *matrix* and for the corresponding map of (split) tori, tensoring with $\mathbb{C}^*$.

Givental's Landau–Ginzburg model is defined by setting the superpotential

$$w = u_1 + \ldots + u_{N+\rho} \in \mathbb{C}[\mathbb{Z}^{N+\rho}],$$

where the u_i are variables corresponding to the basis elements, and restricting to the subvariety defined by $D = 1$, $F_m = 1$ for $m \in [1,l]$, where the F_m are defined by the following:

$$F_m = \sum_{j \in E_m} u_j.$$

Observe that the conditions $D(u_1, \ldots, u_{N+\rho}) = 1$ are monomial, and applying only these equations yields the complex torus $T_{\mathcal{M}} = \operatorname{Spec} \mathbb{C}[\mathcal{N}]$, equipped with the regular function

$$w = \sum_{i \in [1, N+\rho]} z^{r(D_i')},$$

where z^a denotes the monomial in $\mathbb{C}[\mathcal{N}]$ corresponding to $a \in \mathcal{N}$. This is a reformulation of the condition 3.3 in Section 3, setting $q = 1$. Since we have fixed an echelon form for D, $T_{\mathcal{M}}$ has coordinates $\{u_i \mid i \notin E\}$. To pass to the subvariety defined by $\{F_m = 1\}_{m \in [1,l]}$ we introduce new variables y_i for $i \in \bigcup_{m \in [1,l]} E_m^\circ$ defined by the following:

$$u_i = \begin{cases} \frac{1}{1 + \sum_{k \in E_m^\circ} y_k} & \text{if } i = s_m, \\ \frac{y_i}{1 + \sum_{k \in E_m^\circ} y_k} & \text{if } i \in E_m^\circ. \end{cases}$$

So there is a birational torus chart on Givental's Landau–Ginzburg model given by the torus with coordinates $y_j, j \in \bigcup_{m \in [1,l]} E_m^\circ$ and u_j for $j \notin E \cup \bigcup_{m \in [1,l]} E_m$. It is easy to see from our assumptions that w remains Laurent after applying this change of coordinates.

We now apply this procedure explicitly to complete intersections in $P(2, 2+k)$, this is singular, but we recall replacing $\mathrm{Gr}(2, 2+k)$ with $P(2, 2+k)$ is justified in Section 4. Recall that the superpotential $w = \sum_{i \in [1, N+\rho]} z^{r(D_i')}$ for $P(2, 2+k)$, in the coordinates $a_{i,j}$ introduced in Section 4, is given by the Laurent polynomial

$$f_{\mathrm{Gr}(2,2+k)} = a_{1,1} + \sum_{j=2}^{k} \frac{a_{1,j}}{a_{1,j-1}} + \sum_{j=1}^{k} \frac{a_{2,j}}{a_{1,j}} + \sum_{j=2}^{k} \frac{a_{2,j}}{a_{2,j-1}} + \frac{1}{a_{2,k}}.$$

Also consider the $k+2$ equations

$$f_1 = a_{1,1}, \qquad f_j = \frac{a_{1,j}}{a_{1,j-1}} + \frac{a_{2,j}}{a_{2,j-1}}, j \in [2,k], \qquad f_{k+1} = \frac{1}{a_{2,k}}.$$

Given a complete intersection of hypersurfaces of degree d_i, $i \in [1,l]$ we fix a partition

$$[1, k+1] = E_0 \sqcup E_1 \sqcup \ldots \sqcup E_l$$

with $|E_j| = d_j$ for $j > 0$, and we form Givental's Landau–Ginzburg mirror to the complete intersection in $\mathrm{Gr}(2, 2+k)$ by restricting to the subvariety

$$\left\{ F_j = \sum_{r \in E_j} f_r = 1 \mid j = 1, \ldots, l \right\}.$$

In order to apply the method explained above for finding a birational torus chart we apply a change of variables

$$x_{1,1} = a_{1,1}, \qquad x_{1,j} = \frac{a_{1,j}}{a_{1,j-1}}, j \in [2,k],$$
$$x_{2,j} = \frac{a_{2,j}}{a_{2,j-1}}, j \in [2,k], \qquad x_{2,k+1} = \frac{1}{a_{2,k}}.$$

With these changes of variables, of the $3k$ terms of $f_{\mathrm{Gr}(2,2+k)}$, we see that $2k$ are now simply variables $x_{i,j}$ and the remaining k terms are monomials

$$M_i = \frac{a_{2,i}}{a_{1,i}} = \big(\prod_{j \leq i} x_{1,j} \prod_{j \geq i} x_{2,j} \big)^{-1}$$

The polynomials f_j have become the following:

$$f_1 = x_{1,1}, \qquad f_j = x_{1,j} + x_{2,j}, j \in [2,k], \qquad f_{k+1} = x_{2,k+1}.$$

We may now apply the general procedure given above, namely we define the E_m and E used there according to the terms appearing in the polynomials F_m and $\sum_{j=1}^{k} M_j$. Observe that all the requirements for finding a birational torus chart in the first section are met, so we may apply the change of coordinates above. This provides a completely explicit way of forming, potentially several, Laurent polynomial mirrors for a given complete intersection in $\mathrm{Gr}(2, k+2)$. We now consider some examples of this method.

Example A.1 (see also Example 12.12)**.** Consider a cubic intersected with the quadric $\mathrm{Gr}(2,4)$. The weight matrix D for $P(2,4)$ is the following:

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{pmatrix}.$$

The Picard group is the sublattice generated by the vector $(1,1)$. The Laurent polynomial mirror may be obtained either by applying the condition $D = 1$ above, or by changing $f_{\mathrm{Gr}(2,4)}$ to the $x_{i,j}$ variables, in either case there is a Laurent polynomial presentation given by the following:

$$f_{\mathrm{Gr}(2,4)} = x_{1,1} + x_{1,2} + \frac{1}{x_{1,1}x_{1,2}x_{2,3}} + \frac{1}{x_{1,1}x_{2,2}x_{2,3}} + x_{2,2} + x_{2,3}.$$

Where the first k and final k columns of D correspond to basis elements in $\mathcal{N}$ and to variables $x_{1,j}$, $j \in [1,l]$ and $x_{2,j}$, $j \in [2, l+1]$ respectively. The vector $(3,3)$ may be obtained by adding the first, second, fifth, and sixth columns, giving the relation

$$x_{1,1} + x_{1,2} + x_{2,2} + x_{2,3} = 1.$$

Let $E_1 = \{1,2,5,6\}$, $E = \{3,4\}$, and $s_1 = 1$. Denoting the new variables $y_{i,j}$, consistent with the variables $x_{i,j}$ we have the following:

$$x_{1,1} = \frac{1}{1+y_{1,2}+y_{2,2}+y_{2,3}}, \qquad x_{1,2} = \frac{y_{1,2}}{1+y_{1,2}+y_{2,2}+y_{2,3}},$$
$$x_{2,2} = \frac{y_{2,2}}{1+y_{1,2}+y_{2,2}+y_{2,3}}, \qquad x_{2,3} = \frac{y_{2,3}}{1+y_{1,2}+y_{2,2}+y_{2,3}}.$$

The superpotential then becomes

$$\psi^* f_{\mathrm{Gr}(2,4)} = \frac{(1+y_{1,2}+y_{2,2}+y_{2,3})^3}{y_{1,2}y_{2,3}} + \frac{(1+y_{1,2}+y_{2,2}+y_{2,3})^3}{y_{2,2}y_{2,3}} =$$
$$= \frac{(y_{2,2}+y_{1,2})}{y_{1,2}y_{2,2}y_{2,3}}(1+y_{1,2}+y_{2,2}+y_{2,3})^3.$$

We shall now show this is equivalent to the result of Example 12.12 up to symplectomorphisms of cluster type ([CCGGK12], [KP12]). Consider the birational map ϕ_1, defined by the following:

$$\phi_1^* y_{1,2} = y_{1,2}, \qquad \phi_1^* y_{2,2} = y_{2,2}, \qquad \phi_1^* y_{2,3} = (y_{1,2}+y_{2,2})y_{2,3}.$$

Computing $\phi_1^*\psi^* f_{\mathrm{Gr}(2,4)}$ we obtain

$$g_{\mathrm{Gr}(2,4)} = \phi_1^*\psi^* f_{\mathrm{Gr}(2,4)} = \frac{1}{y_{1,2}y_{2,2}y_{2,3}}(1+y_{1,2}+y_{2,2}+(y_{1,2}+y_{2,2})y_{2,3})^3.$$

Now, setting

$$y_{1,2} = a_{2,1}, \qquad y_{2,2} = \frac{a_{2,1}^2}{a_{1,1}}, \qquad y_{2,3} = a_{1,2}$$

we have that

$$g_{\mathrm{Gr}(2,4)} = \frac{a_{1,1}}{a_{2,1}^3 a_{1,2}}\left(1+\left(a_{2,1}+\frac{a_{2,1}^2}{a_{1,1}}\right)(1+a_{1,2})\right)^3 =$$
$$= \frac{a_{1,1}}{a_{1,2}}\left(\frac{1}{a_{2,1}}+\left(1+\frac{a_{2,1}}{a_{1,1}}\right)(1+a_{1,2})\right)^3.$$

Apply another birational change of coordinates ϕ_2, sending

$$a_{i,j} \mapsto a_{i,j}\left(1+\frac{a_{2,1}}{a_{1,1}}\right)^{-1}$$

for each (i,j). We obtain

$$h_{\mathrm{Gr}(2,4)} = \phi_2^* g_{\mathrm{Gr}(2,4)} = \frac{a_{1,1}}{a_{1,2}}\left(\left(1+\frac{1}{a_{2,1}}\right)\left(1+\frac{a_{2,1}}{a_{1,1}}\right)+a_{1,2}\right)^3 =$$
$$= \frac{a_{1,1}}{a_{1,2}}\left(a_{1,2}+\frac{1+a_{1,1}a_{2,1}+a_{1,1}+a_{2,1}}{a_{1,1}a_{2,1}}\right)^3.$$

Which is the resulting polynomial of Example 12.12.

Example A.2 (see also Example 12.6)**.** A fourfold of index 2 given by 4 hyperplane sections of $\mathrm{Gr}(2,6)$. The matrix D for $P(2,6)$ is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The four bundles L_i are all equal to $(1,1,1,1) \in \mathbb{L}^\vee$. We fix the nef-partition by taking the collections of basis elements D_i corresponding to columns $1, 12, \{2,9\}, \{3,10\}$ of the matrix D. Applying the notation employed in Example A.1 we compute

$$\psi^* f_{\mathrm{Gr}(2,6)} = x_{1,4} + x_{2,4} + \frac{(1+y_{2,2})(1+y_{2,3})}{y_{2,2}y_{2,3}x_{2,4}}(1+y_{2,2}+y_{2,2}y_{2,3}) + \frac{(1+y_{2,2})(1+y_{2,3})}{x_{1,4}}.$$

Noting that columns 4, 11 are in neither E nor any of the sets E_m, so the variables $x_{1,4}$ and $x_{2,4}$ persist. As in the previous example, this polynomial also agrees with the result of Example 12.6 up to symplectomorphisms of cluster type, which in particular preserve the period sequence of the Laurent polynomial.

References

[ATY85] L. Aizenberg, A. Tsikh, A. Yuzhakov, *Higher-dimensional residues and their applications Current problems in mathematics*, Encyclopaedia of Mathematical Sciences, Vol. 8, 5–64, 1985.

[Ba97] V. V. Batyrev, *Toric Degenerations of Fano Varieties and Constructing Mirror Manifolds*, Collino, Alberto (ed.) et al., The Fano conference. Papers of the conference, organized to commemorate the 50th anniversary of the death of Gino Fano (1871–1952), Torino, Italy, September 29–October 5, 2002. Torino: Universita di Torino, Dipartimento di Matematica. 109–122 (2004).

[BCFKS97] V. V. Batyrev, I. Ciocan-Fontanine, B. Kim, D. van Straten, *Conifold transitions and mirror symmetry for Calabi–Yau complete intersections in Grassmannians*, Nucl. Phys., B 514, No.3, 640–666 (1998).

[BCFKS98] V. V. Batyrev, I. Ciocan-Fontanine, B. Kim, D. van Straten, *Mirror Symmetry and Toric Degenerations of Partial Flag Manifolds*, Acta Math. 184, No. 1 (2000), 1–39.

[BCFK03] A. Bertram, I. Ciocan-Fontanine, B. Kim, *Two Proofs of a Conjecture of Hori and Vafa*, Duke Math. J. 126, No. 1, 101–136 (2005).

[CCGGK12] T. Coates, A. Corti, S. Galkin, V. Golyshev, A. Kasprzyk, *Mirror Symmetry and Fano Manifolds*, European Congress of Mathematics (Krakow, 2–7 July, 2012), November 2013, pp. 285–300.

[CKP14] T. Coates, A. Kasprzyk, T. Prince, *Four-dimensional Fano toric complete intersections*, arXiv:1409.5030.

[CLS11] D. Cox, J. Little, H. Schenck, *Toric varieties*, Graduate Studies in Mathematics 124. Providence, RI: AMS (2011).

[DH15] C. Doran, A. Harder, *Toric Degenerations and the Laurent polynomials related to Givental's Landau–Ginzburg models*, arXiv:1502.02079.

[DHKLP] C. Doran, A. Harder, L. Katzarkov, J. Lewis, V. Przyjalkowski, *Modularity of Fano threefolds*, in preparation.

[DIM12] O. Debarre, A. Iliev, L. Manivel, *On the period map for prime Fano threefolds of degree 10*, J. Algebr. Geom. 21, No. 1, 21–59 (2012).

[DIM12] O. Debarre, A. Iliev, L. Manivel, *Special prime Fano fourfolds of degree 10 and index 2*, arXiv:1302.1398.

[EHX97] T. Eguchi, K. Hori, C.-Sh. Xiong, *Gravitational quantum cohomology*, Int. J. Mod. Phys. A 12, No. 9, 1743–1782 (1997).

[Ga07] S. Galkin, *Small toric degenerations of Fano 3-folds*, preprint `http://www.mi.ras.ru/~galkin/work/3a.pdf`.

[Gi97a] A. Givental, *Stationary phase integrals, quantum Toda lattices, flag manifolds and the mirror conjecture*, Topics in singularity theory, 103–115, Amer. Math. Soc. Transl. Ser. 2, 180, AMS, Providence, RI, 1997.

[Gi97b] A. Givental, *A mirror theorem for toric complete intersections*, Topological field theory, primitive forms and related topics (Kyoto, 1996), 141–175, Progr. Math., 160, Birkhauser Boston, Boston, MA, 1998.

[GKR12] M. Gross, L. Katzarkov, H. Ruddat, *Towards Mirror Symmetry for Varieties of General Type*, arXiv:1202.4042.

[HV00] K. Hori, C. Vafa, *Mirror symmetry*, arXiv:hep-th/0002222.

[ILP13] N. Ilten, J. Lewis, V. Przyjalkowski. *Toric Degenerations of Fano Threefolds Giving Weak Landau–Ginzburg Models*, Journal of Algebra 374 (2013), 104–121.

[IP99] V. Iskovskikh, Yu. Prokhorov, *Fano varieties*, Encyclopaedia of Mathematical Sciences, **47** (1999) Springer, Berlin.

[Ki00] B. Kim, *Quantum hyperplane section principle for concavex decomposable vector bundles*, J. Korean Math. Soc. 37 (2000), no. 3, 455–461.

[Ko94] M. Kontsevich, *Homological algebra of mirror symmetry*, Proc. International Congress of Matematicians (Zürich 1994), Birkhäuzer, Basel, 1995, pp. 120–139.

[KP12] L. Katzarkov, V. Przyjalkowski, *Landau–Ginzburg models — old and new.* Akbulut, Selman (ed.) et al., Proceedings of the 18th Gokova geometry–topology conference. Somerville, MA: International Press; Gokova: Gokova Geometry-Topology Conferences, 97–124 (2012).

[Lee01] Y. Lee, *Quantum Lefschetz hyperplane theorem*, Invent. Math. 145 (2001), no. 1, 121–149.

[Li13] Zh. Li, *On the birationality of complete intersections associated to nef-partitions*, arXiv:1310.2310.

[Ma99] Yu. Manin, *Frobenius manifolds, quantum cohomology, and moduli spaces*, Colloquium Publications. American Mathematical Society (AMS). 47. Providence, RI: American Mathematical Society (AMS) (1999).

[MR13] R. Marsh, K. Rietsch, *The B-model connection and mirror symmetry for Grassmannians*, arXiv:1307.1085.

[Prz08] V. Przyjalkowski, *On Landau–Ginzburg models for Fano varieties*, Comm. Num. Th. Phys., Vol. 1, No. 4, 713–728 (2008).

[Prz10] V. Przyjalkowski, *Hori–Vafa mirror models for complete intersections in weighted projective spaces and weak Landau–Ginzburg models*, Cent. Eur. J. Math. 9, No. 5, 972–977 (2011).

[Prz13] V. Przyjalkowski, *Weak Landau–Ginzburg models for smooth Fano threefolds*, Izv. Math. Vol., 77 No. 4 (2013), 135–160.

[PSh14] V. Przyjalkowski, C. Shramov, *On weak Landau–Ginzburg models for complete intersections in Grassmannians*, Russian Math. Surveys 69, No. 6, 1129–1131 (2014).

[PSh15a] V. Przyjalkowski, C. Shramov, *On Hodge numbers of complete intersections and Landau–Ginzburg models*, IMRN, 2015, doi: 10.1093/imrn/rnv024.

[PSh15b] V. Przyjalkowski, C. Shramov, *Laurent phenomenon for Landau–Ginzburg models of complete intersections in Grassmannians*, Proc. Steklov Inst. Math. 290 (2015).

[St93] B. Sturmfels, *Algorithms in Invariant Theory*, Texts and Monographs in Symbolic Computation, Wien, Springer–Verlag, 1993.

Steklov Mathematical Institute of Russian Academy of Sciences, 8 Gubkina st., Moscow 119991, Russia

E-mail address: `victorprz@mi.ras.ru, costya.shramov@gmail.com`